

A List of Known Ant Species of Thailand (Formicidae: Hymenoptera)

WEEYAWAT JAITRONG^{1*} AND JARUJIN NABHITABHATA¹

¹ Thailand Natural History Museum, National Science Museum, Technopolis, Khlong 5, Khlong Luang,
Pathum Thani 12120, THAILAND

ABSTRACT - The presently 247 known ant species were recorded to occur in Thailand and distributing among 55 genera in 9 subfamilies. Six of them were originally described directly from Thai territory, i.e. *Aenictus artipus*, *Leptanilla thai*, *Lophomyrmex striatulus*, *Tetramorium ciliatum*, *Tetramorium flavipes*, and *Tetraponera notabilis*. Information provided for each name include the valid scientific name and its author and year, reference to the original designation, the type locality and known distribution of each particular species in Thailand.

KEY WORDS : Ants, Check list, Thailand, Distribution.

INTRODUCTION

Ants are well-recognized eusocial insects belong to the family Formicidae within the order Hymenoptera the same with bees and wasps. It is well known that they can be found in almost all terrestrial habitats on earth, from cold subarctic tundra to moist equatorial rain forest, from waterlogged swamp to harsh desert, from sea coast to high altitudes, and from deep in the soil to the tips of highest trees (Bolton, 1994). They are usually soil dwellers and acting the important role in the terrestrial ecosystem as controllers of the numbers of other insects or small soil invertebrates.

There are approximately 15,000 living ant species, of which some 9,000 - 10,000 species have been described and classified into 16 subfamily in the world (Bolton, 1994). At present, 9 subfamilies of ants are recorded in Southeast Asia and all found to be

native to Thailand with the estimated number of 800-1000 species.

In recent years around the world interests on ants had inspired the production of large number of reports concerning the varied aspects of biology and behavior. Dealing with the ant faunal surveys that antedated 1950, they are recommended not to use in the present studies, as their provided keys are clogging with unavailable names (Bolton, 1995). In the period between 1950-2003 have seen the appearance of the number of ant surveys locally that normally including keys for the identification of genera and species found within such country or territory, and also lists or catalogues of existing ants provided.

Beginning as lately as 1906, the first reliable records of ant faunas of Thailand appeared on the southern ant species (Bingham, 1906), where he discovered 39 species from Songkhla, Pattani and Narathiwat. Until recently our ant fauna has been regarded as poorly known, comparing to those of the neighboring countries. Those entomologists who had contributed, more or less, to the knowledge of the Thai ants include Emery (1893c), Wilson

* Corresponding author.
Tel. (662) 577 9999 ext. 1506
E-mail: polyrhachis@yahoo.com

(1964), Baroni (1977), Bolton (1977), Rigato (1994) and Ward (2001). For Thai myrmecologists, they mostly studied on the biological diversity of ants for example, Watanasit et al. (2000 and 2003) reported on the diversity of ants in Songkhla Province and their preliminary survey of ant in Satul Province, both in Southern Thailand; only one the taxonomic study of Thai ants was done by Wiwatwitaya and Jaitrong (2001) on ants in Khao Yai National Park in the north-eastern Thailand.

MATERIALS AND METHODS

Starting in period 1997 - 2004, the taxonomic surveys of ant faunas in various localities throughout the country had been conducted simultaneously with the review works on Thai ant species reported or described as new taxa in revisionary works on species groups or records of native species of nearby countries in the region.

Entries are arranged alphabetically with the accepted valid name of genera, and within each genus, of species. Name of subfamily and tribe is in parenthesis after the generic name. The generic name is provided with author and year of establishment, the original citation; the type species and type designation of the scientific name, also indicated, the author and year, the original citation and the type locality. Distribution of all known collecting localities are shown by regional part; starting from the north, to northeast, east, southeast, centre, southwest, lastly to the peninsular. The province shown in italic type is followed by the name of the national park, wildlife sanctuary, or district.

This list is here regarded as a preliminary one. As only 10 % of the country have been intensively surveyed for their respective ant faunas. So large extense of land still await further discovery.

Abbreviations used.- The following abbreviations will be found appearing regularly in the list:

- (s.) soldier caste or major worker, when appropriate.
(w.) worker.

- (q.) queen.
(m.) male.
IS. Island.
subsp. subspecies
var. variety

RESULTS

A LIST OF ANT SPECIES

ACANTHOMYRMEX (Myrmicinae: Myrmicini)

สกุลมดหัวดิน

Acanthomyrmex Emery, 1893a: cclxxvi. *Bulletin Bimensuel de la Societe Entomologique de France 1892* No. 20 cclxxv-cclxxvii. Type species: *Acanthomyrmex luciolae* Emery, 1893c, by subsequent designation of Bingham, 1903: 191.

Acanthomyrmex ferox Emery, 1893c

มดหัวดินมลายู

Acanthomyrmex ferox Emery, 1893c: *Annales de la Societe Entomologique de France* 62: 245, pl. 6, fig.11. (w). Type locality: West Malaysia.

DISTRIBUTION :

PENINSULAR: *Ranong*: Klong Yan; *Nakhon Si Thammarat*: Khao Luang; *Narathiwat*: Halabala.

ACROPYGA (Formicinae: Plagilepidini)

สกุลมดหัวละม้ายท้าย

Acropyga Roger, 1862: 242. *Berliner Entomologische Zeitschrift* 6: 233-254. Type species: *Acropyga acutiventris* Roger, 1862, by monotypy.

Acropyga acutiventris Roger, 1862

มดหัวละม้ายท้ายใหญ่

Acropyga acutiventris Roger, 1862: *Berliner Entomologische Zeitschrift* 6: 243. (w.q.) Type locality: Sri Lanka.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Inthanon, Doi Suthep-Pui, Doi Chiang Dao, Mae Tang, Mae Cham; *Chiang Rai*: Mae Pha Luang; *Lampang*: Tham Pha Thai; *Tak*: Thung Yai.
EASTERN: *Nakhon Ratchasima*: Khao Yai,

Sakaerat; *Ubon Ratchathani*:
Pha Taem.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu
Nai; *Chon Buri*: Siracha.

PENINSULAR: *Chumphon*: Krom Luang Chumphon;
Ranong: Klong Naka, Klong Yan;
Nakhon Si Thammarat: Khao
Luang; Trang: Khao Cho *Narathiwat*:
Toa Dang, Halabala.

***AENICTUS* (Aenictinae)**

สกุลมดทหาร

Aenictus Shuckard, 1840: 266. *Annals of
Natural History* 5: 258-271. Type species: *Aenictus
ambiguus* Shuckard, 1840, by original designation.

***Aenictus artipus* Wilson, 1964**

มดทหารสยาม

Aenictus artipus Wilson, 1964: *Pacific
Insects* 6 (3): 449, fig. 60 (w). Type locality: Chiang
Mai, Thailand.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Inthanon.
EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon
Ratchasima*: Khao Yai.

***Aenictus binghami* Forel, 1900a**

มดทหารบิงแฮม

Aenictus binghami Forel, 1900a: *Annales
de la Societe Entomologique de Belgique* 44: 76(w).
Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Ang Khang,
Doi Chiang Dao; *Phrae*: Wang
Chin; *Tak*: Thung Yai; *Phisanulok*:
Thung Salaeng Luang.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon
Ratchasima*: Sakaerat, Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu
Nai; *Chanthaburi*: Khao Soi
Dao, Nam Tok Phliu.

PENINSULAR: Trang: Khao Chong; *Narathiwat*:
Halabala.

***Aenictus camposi* Wheeler and Chapman, 1925**

มดทหารเตลียง

Aenictus camposi Wheeler and Chapman,
1925: *Philippine Journal of Science* 28: 48, pl. 1, figs.

3, 4. (w). Type locality: Philippines.

DISTRIBUTION:

NORTHERN: *Phitsanulok*: Thung Salaeng Luang.

EASTERN: *Nakhon Ratchasima*: Khao Yai,
Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu
Nai; *Chanthaburi*: Khao Soi
Dao.

***Aenictus ceylonicus* (Mayr, 1866a)**

มดทหารศรีลังกา

Typhlatta ceylonicus Mayr, 1866a:
*Sitzungsberichte der k. Akademie der Wissenschaften.
Mathematisch Naturwissenschaftliche Classe* 53: 505.
(w). Type locality: Sri Lanka.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon
Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Aenictus dentatus* Forel, 1911c**

มดทหารทราย

Aenictus aitikeni var. *dentatus* Forel, 1911c:
Bulletin de la Societe Vaudoise des sciences Naturelles
47: 383. (w). Type locality: West Malaysia.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae
Tang; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon
Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang
Reu Nai.

PENINSULA: *Narathiwat*: Halabala.

***Aenictus fergusonii* Forel, 1901a**

มดทหารอกชน

Aenictus fergusonii Forel, 1901a: *Journal of
the Bombay Natural History Society* 13: 473. (w). Type
locality: India.

DISTRIBUTION:

NORTHERN: Chiang Mai: Doi Ang Khang.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Aenictus hottai* Terayama and Yamane, 1989**

มดทหารมลายู

Aenictus hottai Terayama and Yamane,
1989: *Japanese Journal of Entomology* 57: 598, figs.

1, 2. (w). Type locality: Sumatra.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

***Aenictus javanus* Mayr, 1896**

มดทหารขวา

Aenictus javanus Mayr, 1896: *Annales de la Societe Entomologique de Belgique* 40: 245. (m). Type locality: Java.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

CENTRAL: *Saraburi*: Phu Kae.

***Aenictus laeviceps* (F. Smith, 1857)**

มดทหารอกรเขียว

Typhlatta laeviceps F. Smith, 1857: *Journal of Proceedings of the Linnean Society of London, Zoology* 2: 79. (w). Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Ang Khang;

Tak: Thung Yai; *Phitsanulok*: Thung Salaeng Luang.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Narathiwat*: Halabala.

***AMBLYOPONE* (Ponerinae: Amblyoponini)**

สกุลมดปากเลื้อย

Amblyopone Erichson, 1842 : 260. *Archiv fur Naturgeschichte* 8 : 83-287. Type species: *Amblyopone australis* Erichson, 1842, by monotypy. *Amblyopone* senior synonym of *Stigmatomma*: Emery and Forel, 1879 : 455; Mayr, 1887 : 546.

***Amblyopone reclinata* Mayr, 1879**

มดปากเลื้อยดำใหญ่

Amblyopone reclinata Mayr, 1879 : *Verhandlungen der k.k. Zoologisch- Botanischen Gesellschaft in Wien* 28. 667. (w). Type locality: West Java.

DISTRIBUTION:

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat

CENTRAL: *Saraburi*: Phu Kae.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Trang*: Khao Chong.

***ANOCHEBUS* (Ponerinae: Ponerini)**

สกุลมดลั่นไก

Anochetus Mayr, 1861: 53. *Die Europaischen Formiciden*: 80 pp. Type species: *Odontomachus ghilianii* Spinola, 1851, by monotypy. *Anochetus senior* synonym of *Stenomymex*: Forel, 1887: 382. *Anochetus senior* synonym of *Myrmapetes*: Brown, 1953b: 2. *Anochetus* provisional junior synonym of *Odontomachus*: Brown, 1973: 178.

***Anochetus graeffei* Mayr, 1870**

มดลั่นไกธรรมดา

Anochetus graeffei Mayr, 1870: *Verhandlungen der k.k. Zoologisch- Botanischen Gesellschaft in Wien* 2: 961. (w). Type locality: Samoa.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao, Doi Ang Khang, Ob Luang, Doi Luang, Doi Suthep-Pui, Doi Inthanon, Mae Tang, Mae Cham; *Lampang*: Thum Pha Thai, Hai Tak teak plantation; *Phrae*: Wang Chin; *Tak*: Thung Yai, Lan Sang

NORTHEASTERN: *Kalasin*: Phu Si Than.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat; *Ubon Ratchathani*: Pha Taem.

SOUTHWESTERN: *Kanchanaburi*: Thong Pha Phum, Hai Kha Khang.

CENTRAL: *Saraburi*: Phu Kae; *Bangkok*: Bang Khen, Chatuchak.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chon Buri*: Siracha; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Chumphon*: Krom Luang Chumphon; *Ranong*: Klong Naka, Klong Yan; *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Songkhla*: Ton Nga Chang, Khao Kor Hong; *Narathiwat*: Halabala, Toa Dang.

***Anochetus modicus* Brown, 1978**

มดลินไก่อะโหลกเหลือง

Anochetus modicus Brown, 1978: 582. *Studia Entomologica* (N.S.) 20: (w, q, m). Type locality: Borneo.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Anochetus myops* Emery, 1893b**

มดลินไก่อะโหลกเล็ก

Anochetus myops Emery, 1893b: 20. *Revue Suisse de Zoologie* 1: 201, pl. 8, figs 11, 12. (w). Type locality: West Malaysia.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Anochetus princeps* Emery, 1884**

มดลินไก่อานม้า

Anochetus princeps Emery, 1884. *Annali del Museo Civico di Storia Naturale di Genova* (2) [27]: 379. (Diagnosis in Key) (w). Type locality: Java.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Anochetus rugosus* (F. Smith, 1857)**

มดลินไก่ออกแหลม

Odontomachus rugosus F. Smith, 1857. *Journal of Proceedings of the Linnean Society of London, Zoology* 2: 65 (w) Type locality: Singapore.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***ANOPOLEPIS* (Formicinae: Plagiolepidini)**

สกุลมดน้ำผึ้ง

Anoplolepis Santschi, 1914. 123 [as subgenus of *Plagiolepis*]. *Insectes Hymenopteres* 2. Formicidae: 41-148. Paris. Type species: *Formica longipes* F. Smith, 1857. 55 (w.) (junior primary homonym in *Formica*, *Anoplolepis gracilipes* first available replacement name), by monotypy.

***Anoplolepis gracilipes* (F. Smith, 1857)**

มดน้ำผึ้ง

Formica gracilipes F. Smith, 1857. *Journal of Proceedings of the Linnean Society of*

London, Zoology 2: 73. (w) Type locality: Singapore.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao, Doi Ang Khang, Ob Luang, Doi Luang, Doi Suthep-Pui, Doi Inthanon, Mae Tang, Mae Cham; *Lampang*: Tham Pha Thai, Huai Tak teak plantation; *Phrae*: Wang Chin; Tak: Thung Yai, Lan Sang

NORTHEASTERN: *Kalasin*: Phu Si Than.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat; *Ubon Ratchathani*: Pha Taem.

CENTRAL: *Saraburi*: Phu Kae; *Bangkok*: Bang Khen, Chatuchak.

SOUTHWESTERN: *Kanchanaburi*: Thong Pha Phum, Huai Kha Khang.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chon Buri*: Siracha; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Chumphon*: Krom Luang Chumporn; *Ranong*: Klong Naka, Klong Yan; *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Songkhla*: Ton Nga Chang, Khao Kor Hong; *Narathiwat*: Halabala, Toa Dang.

***CALYPTOMYRMEX* (Myrmicinae: Stenammini)**

สกุลมดปากล้อม

Calyptomyrmex Emery, 1887b: 471. *Annali del Museo Civico di Storia Naturale di Genova* (2) 5: 528-534. Type species: *Calyptomyrmex beccarii* Emery, 1887b, by monotypy. *Calyptomyrmex* senior synonym of *Weberdris*: Donisthorpe, 1949: 84. Revision of Oriental species: Baroni, 1975a: 410; of Afrotropical species: Bolton, 1981: 61.

***Calyptomyrmex emeryi* Forel, 1901b**

มดปากล้อม

Calyptomyrmex emeryi Forel, 1901b: *Jahrbuch der Hamburgischen Wissenschaftlichen Anstalten [Mitteilungen aus dem Naturhistorischem Museum]* 18: 51. (w.m). Type locality: Borneo.

DISTRIBUTION:PENINSULAR: *Narathiwat*: Halabala.**CAMPONOTUS (Formicinae: Camponotini)****สกุลมดตะลัน**

Camponotus Mayr, 1861: 35. *Die Europaischen Formicidem in Wein*: 80 pp. Type species: *Formica ligniperda* Latreille, 1802: 88, by subsequent designation of Bingham, 1903: 347. Subgenera of *Camponotus*: *Colobopsis*, *Condylomyrma* (junior synonyms of *Colobopsis*), *Dinomyrmex*, *Hypercolobopsis*, *Karavaievia*, *Manniella*, *Mayria*, *Myrmacrhaphe*, *Myrmemblys*, *Myrmapheanus*, *Myrmentoma*, *Myrmepinotus*, *Myrmepomis*, *Myrmespera*, *Myrmeurynota*, *Myrmisolepis*, *Myrmobrachys*, *Myrmocamelus*, (junior synonyms of *Myrmophyma*), *Myrmocladoecus*, *Myrmodirachis*, *Myrmogigas*, (junior synonyms of *Dinomyrmex*), *Myrmogonia*, *Myrmolophus* (junior synonyms of *Myrmepomis*), *Myrmomalis*, *Myrmonesites*, *Myrmopalpella*, *Myrmopelta*, *Myrmophyma*, *Myrmopiromis*, *Myrmoplatypus*, *Myrmoplatys*, *Myrmopsamma*, *Myrmopytia*, *Myrmosaga*, *Myrmosaulus*, *Myrmosericus*, *Myrmospincta*, *Myrmostenus*, *Myrmotarsus*, *Myrmotemnus*, *Myrmothrix*, *Myrmotrema*, *Myrmoturbra* (junior synonym of *Tanaemyrmex*), *Myrmoxxygenys*, *Neocolobopsis* (junior synonym of *Hypercolobopsis*), *Neomyrmemblys* (junior synonym of *Myrmapheanus*), *Orthonotomyrmex*, *Orthonotus*, *Paracolobopsis*, *Paramyrmemblys*, *Pseudocolobopsis*, *Rhinomyrmex*, *Tanaemyrmex*, *Thlipsepinotus*. Synopses of genera: Wheeler, W. M. 1922:703; Emery, 1925: 62. All of genera given as provisional junior synonym of *Campouts*: Brown, 1973: 179-185.

Camponotus arrogans* (F. Smith, 1858)*มดตะลันขนอนุกรมกัมมัน**

Formica arrogans F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum 6 Formicidae*: 23 (w). Type locality: Singapore.

DISTRIBUTION:SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu NaiPENINSULAR: *Songkhla*: Ton Nga Chang;*Narathiwat*: Halabala.***Camponotus auriventris* Emery, 1889****มดตะลันพม่า**

Camponotus auriventris Emery, 1889: *Annali del Museo Civico di Storia Naturale di Genova* (2) 7: 514 (s.w.q). Type locality: Myanmar.

DISTRIBUTION:NORTHERN: *Chiang Mai*: Mae Cham; *Tak*: Thung Yai.EASTERN: *Nakhon Ratchasima*: Khao Yai.SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.PENINSULAR: *Ranong*: Klong Naka***Camponotus camelinus* (F. Smith, 1857)****มดตะลันป่าหัวดำ**

Formica camelinus F. Smith, 1857: *Journal of proceeding of the Linnean Society of London, Zoology* 2:57 (w). Type locality: Singapore.

DISTRIBUTION:NORTHERN: *Chiang Mai*: Mae Cham, Doi Chiang Dao; *Lampang*: Tham Pha Thai; *Tak*: Thung Yai.EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.PENINSULAR: *Trang*: Khao Chong; *Narathiwat*: Halabala, Toa Dang.***Camponotus cylindricus* (Fabricius, 1798)****มดตะลันดำหัวแดง**

Formica cylindrica Fabricius, 1798: *Supplementum Entomologiae Systematicae*: 280. (w). Type Locality: Borneo.

DISTRIBUTION:PENINSULAR: *Ranong*: Klong Naka***Camponotus dolichoderoides* Forel, 1911a****มดตะลันกันห้อย**

Camponotus dolichoderoides Forel, 1911a: *Revue Suisse Zoologie* 19: 51, (w). Type locality: Borneo

DISTRIBUTION:PENINSULAR: *Narathiwat*: Halabala.***Camponotus gigas* (Latreille, 1802)****มดตะลันยักษ์ปีกขี้ไต้**

Formica gigas Latreille, 1802: *Histoire Naturelle des Fourmis*, 105 pl. 2, fig.6 (s). Type locality: Grandes-Indes. Subgenus *Dinomyrmex*.

DISTRIBUTION:

PENINSULAR: *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala, Toa Dang.

Camponotus lasiselene* Wang and Wu, 1994*มดตะลานยอดไม้**

Camponotus lasiselene Wang and Wu, 1994: *Journal of Beijing Forestry University (English edition)* 3. 24, fig. 3 (w). Type locality: China.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui; *Lampang*: Tham Pha Thai, Huai Tak; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

CENTRAL: *Bangkok*: BangKhen.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chon Buri*: Siracha; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

Camponotus leonadi* Emery, 1889*มดตะลานขนทอง**

Camponotus (Colobopsis) leonadi Emery, 1889: *Annali del Museo Civico di Storia Naturale di Genova* (2) 7. 515, pl.11, figs. 22, 23 (s.w.). Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Ob Luang, Mae Tang, Doi Suthep-Pui; *Lampang*: Tham Pha Thai, Mae Had, Ngao; *Phrae*: Wang Chin; *Tak*: Thung Yai; *Phitsanulok*: Thung Salaeng Luang.

NORTHEASTERN: Kalasin: Phu Si Than.

EASTERN: *Nakhon Ratchasima*: Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Trang*: Khao Chong; *Songkhla*: Khao Nam Khang; *Narathiwat*: Halabala, Toa Dang.

Camponotus nicobarensis* Mayr, 1865*มดตะลานนิโคบาร์**

Camponotus nicobarensis Mayr, 1865: *Zoologischer Theil. Formicidae*. 31, pl. 1, fig. 1(w) Type locality: Nicobaris

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Trang*: Khao Chong; *Narathiwat*: Halabala, Toa Dang

Camponotus rufifemur* Emery, 1900*มดตะลานขาขน**

Camponotus rufifemur Emery, 1900: *Annali del Museo Civico di Storia Naturale di Genova* (2) 20. 705 (s.q.). Type locality: Indonesia

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Toa Dang, Halabala.

Camponotus rufoglaucus* (Jerdon, 1851)*มดตะลานปล้องขี้เถ้า**

Formica rufoglaucus Jerdon, 1851: *Madras Journal of Literature and Science* 17: 124 (w). Type locality: India.

DISTRIBUTION:

Found in all provinces.

Camponotus saundersi* Emery, 1889*มดตะลานแดง**

Camponotus (Colobopsis) saundersi Emery, 1889: *Annali del Museo Civico di Storia Naturale di Genova* (2) 7. 516 (s.w.). Type locality: Myanmar.

DISTRIBUTION:

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala, Toa Dang.

Camponotus sericeus* (Fabricius, 1798)*มดตะลานเสี**

Formica cericea Fabricius, 1798: *Supplementum Entomologiae Systematical*: 279, (w.). Type locality: Senegal.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui; *Lampang*: Tham Pha Thai, Ngao.

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat.

Camponotus singularis* (F. Smith, 1858)*มดตะลันป่าหัวแดง**

Formica singularis F. Smith, 1858: *Catalogue of Hymenopteous Insects in The collection of British Museum, 6 London*. 27 (w.). Type locality: Java.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

PENINSULAR: *Narathiwat*: Halabala.

Camponotus vitreus* (F. Smith, 1860b)*มดตะลันออกคอดดำ**

Formica vitrea F. Smith, 1860b: *Journal of the Proceedings of the Linnean Society, Zoology* 5 (4). 94 (w.). Type locality: Indonesia.

DISTRIBUTION:

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Songkhla*: Ton Nga Chang.

CARDIOCONDYLA* (Myrmicinae: Formicoxenini)*สกุลมดเกล็ดเรียบ**

Cardiocondyla Emery, 1869: 20. Type species: *Cardiocondyla elegans* Emery, 1869. by monotype. *Cardiocondyla* senior synonym of *Emeryia*: Forel, 1892a: 461; Foneil, 1892b: 313. *Cardiocondyla* senior synonym of *Xenometa*: Baroni Urbani, 1973: 200. *Cardiocondyla* senior synonyms of *Dyclona*, *Loncyda* and *Prosopidris*: Bolton, 1982: 309; Revision of Palaearctic species: Bernard, 1956: 299; Afrotropical species: Bolton, 1982, 312.

Cardiocondyla emeryi* Forel, 1881*มดเกล็ดเรียบอีเมอริ**

Cardiocondyla emeryi Forel, 1881: *Annali del Museo Civico di Storia Naturale di Genova* 16: 5, (w.). Type locality: Virginis.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang.

EASTERN: *Nakhon Ratchasima*: Sakaerat, Khao Yai.

CENTRAL: *Bangkok*: Kasetsart University.

PENINSULAR: *Trang*: Khao Chong; *Songkhla*: Khao Nam Khang; *Narathiwat*: Halabala.

Cardiocondyla nuda* (Mayr, 1866a)*มดเกล็ดเรียบลีตาด**

Leptothorax nudus Mayr, 1866a: *Mathematisch-Naturwissenschaftliche Class* 53. 508 (w.). Type locality: Fijlis.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang, Ob Luang, Doi Suthep-Pui.

EASTERN: *Nakhon Ratchasima*: Sakaerat, Kho Yai.

PENINSULAR: *Songkhla*: Khao Nam Khang.

Cardiocondyla wroughtonii* (Forel, 1890)*มดเกล็ดเรียบเหลือง**

Emeryia wroughtonii Forel, 1890: *Annales de la Societe Entomologique de Belgique. Comptes-rendus* 34. Cxi, (ergatoid m., not w.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang, Mae Cham, Doi Suthep-Pui.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

PENINSULAR: *Songkhla*: Khao Nam Khang.

CAREBARA* (Myrmicinae: Pheidologetonini)*สกุลแมงมัน**

Carebara Westwood, 1840: 86. Type species: *Carabara lignata* Westwood, 2840, by monotype. Key to Afrotropical species: Wheeler, W.M. 1922a 172. Review of genus: Ettershank, 1966:125.

Carebara castanea* F. Smith, 1858*แมงมัน**

Carebara castanea F. Smith, 1858: *Catalogue of Hymenopterous Insects in collection of British Museum* 6 Formicidae: 178 (q.) Type locality: Hong Kong

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang, Phang, Doi Suthep-Pui, Doi Chiang Dao; *Phayao*: Mae Ka; *Lampang*: Tham Pha Thai, Ngao.

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat.

CATAULACUS (Myrmicinae: Cataulacini)**สกุลมดปล้องไฟ**

Cataulacus F. Smith, 1853: 225. Type species: *Cataulacus taprobanea* F. Smith, 1853, by subsequent designation of *Otomymex* and Revision of genus: Botton, 1974a: 1. Key to Afrotropical species: Botton, 1982: 354.

Cataulacus granulatus* (Latreille, 1802)*มดปล้องไฟธรรมดา**

Formicia granulatus Latreille, 1802: *Histoire Naturelle des forems, et recueil dememoriees et d(observations sur les abeilles, Les araignees, les faucheurs, et autres insectes*: 445 pp. Paris. 275, pl. 12, fig. 75 (w.) Type locality: Grand Indis.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang, Doi Suthep-Pui, Doi Inthanon, Ob Luang; *Chiang Rai*: Mae Pha luang; *Phayao*: Mae Yluar; *Phrae*: Wang Chin; *Tak*: Thung Yai.

NORTHEATERN: *Kalasin*: Phu Si Than.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHWESTERN: *Uthai Thani*: Huai Kha Kang; *Kanchanaburi*: Thong Pha Phum; *Prachaup Khiri Khan*: Huai Yang.

CENTRAL: *Saraburi*: Phu Kae; *Bangkok*: Bang Khen, Kesetsart University.

SOUTHEASTERN: *Sa Kaeo*: Nam Tok Khao Takrub; *Prachin Buri*: Khao Yai; *Chachoengsao*: Khao Aang Reu Nai; *Chon Buri*: Siracha; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu; *Trat*: Ko Chang.

PENINSULAR: *Chumphon*: Krom Luang Chumphon; *Ranong*: Klong Yan, Klong Naka. *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong, Palian; *Songkhla*: Khoa Khor Hong, Ton Nga Chang, Khao Nam Khang, Hat Yai; *Narathiwat*: Halabala, Toa Dang.

Cataulacus horridus* F. Smith, 1857*มดปล้องไฟปีกมิได้**

Cataulacus horridus F. Smith, 1857: *Journal of the Proceedings of the Linnean Society of London, Zoology* 2: 81, pl.2, fig.3 (w.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

CENTROMYRMEX (Ponerinae: Ponerini)**สกุลมดตีนหนาม**

Centromyrmex Mayr, 1866b: 894. Type species: *Centromyrmex bohemani* (junior synonym of *Centromyrmex brachycola*, by monotypy. *Centromyrmex* senior synonym of *Spalacomymex*: Emery, 1890: 40. *Centromyrmex* senior synonym of *Typhloteras*: Brown, 1953a: 8. *Centromyrmex* senior synonym of *Glyphopone* and *Leptopone*: Brown 1963: 9. *Centromyrmex* senior of synonym of *Promyopias*: Botton, 1994: 194. Revision of Neotropical species: Kempf, 1967: 401.

Centromyrmex feae* (Emery, 1889)*มดตีนหนามแดง**

Spalacomymex feae Emery, 1889: *Annali del Museo Civio di Storia Naturale di Genova* (2) 7. 491, pl. 10, figs. 11-15 (w.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang, Doi Chiang Dao; *Phayao*: Mae Ka, Mae Yluar; *Lampang*: Tham Pha Thai; *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat.

CENTRAL: *Saraburi*: Phu Kae.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu; *Trat*: Ko Chang.

PENINSULAR: *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala.

CERAPACHYS (Cerapachyinae: Cerapachyini)**สกุลมดท้องคอด**

Cerapachys F. Smith, 1857: 74. Type

species: *Cerapachys antennatus* Smith, F. 1857, by subsequent designation of Bingham, 1903. 28. *Cerapachys* senior synonym of *Cerapachys*: Wheeler, W. M. 1922: 755. *Cerapachys* senior synonym of *Parasyscia* and *Syscia* (and its junior synonym *Cyscia*): Kempf, 1972: 76. *Cerapachys* senior *Chrysapace*, *Lioponera*, *Neophyracaces*, *Ooceraea*, *Phyracaces*, *Procerapachys*, and Revision of genus: Brown, 1975:18

***Cerapachys dohertyi*. Emery, 1902**

มดท้องคอดหูยาว

Cerapachys doherityi. Emery, 1902: Rendiconto della Sessioni delle R. Accademia delle Scienze dell(Istituto di Bologna (N.S.) 6. 25, (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

***Cerapachys longitarsus* (Mayr, 1879)**

มดท้องคอดตีนยาว

Lioponera longitarsus Mayr, 1879: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 28. 677(w.q.) Type locality: India.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.

***Cerapachys sexspenus* (Xu, 2000)**

มดท้องคอดยูนาน

Yunodorylus sexspenus Xu, 2000: *Zoological Research* 4: 298, figs. 1-6 (w.) Type locality: China.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham; *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Sakaerat.

***Cerapachys sulcinodis*, Emery, 1889**

มดท้องคอดสีนิล

Cerapachys sulcinodis Emery, 1889: *Annali del Museo Civico di Storia Naturale di Genova* (2) 7. 493 (w.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang, Doi Suthep-Pui, Doi Chiang Dao; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Trang*: Khao Chong; *Narathiwat*: Halabala.

CREMATOGASTER (Myrmicinae: Crematogastrini)

สกุลมดอี (หนือ) หรือ มดสอด (ใต้)

Crematogaster Lund, 1831. 132: Type species: *Formica scutellaris* Oliver, 1792, by subsequent designation of Bingham, 1903:124. [Type species not *Formica acuta*, unjustified subsequent designation by Emery, 1912, 272; this error repeated in, for example Emery, 1922: 128.] *Crematogaster* senior synonym of *Acrocoelia* Mayr, 1863:404 (see also Buren, 1959: 125, Kempf, 1972: 81). *Crematogaster* senior synonym of *Tranopeltoides*: Kempf, 1960. 173. [*Crematogaster* Mayr, 1861: 74 (and many later authors), incorrect subsequent spelling. Discussion of spelling: Emery, 1912: 272 (foot note); Wheeler., W.M., 1913a: 78; Donithorpe, 1941c: 36 and Buren, 1959 subsequent of *Crematogaster*: nominal plus *Apteroкрема*, *Atopogyne*, *Colobocrema*, *Decacrema*, *Eucrema*, *Mesocrema*, *Nematocrema*, *Neocrema*, *Orthocrema*, *Oxygyne*, *Paracrema*, *Physocrema*, *Sphaerocrema*, *Xiphocrema*.

***Crematogaster coriaria* Mayr, 1872**

มดอีดำ

Crematogaster coriaria Mayr, 1872: *Annali del Museo Civico di Storia Naturale di Genova* 2: 154 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Trang*: Khao Chong; *Songkla*: Ton Nga Chang.

***Crematogaster difformis* F. Smith, 1857**

มดอีโป่งดำ

Crematogaster difformis F. Smith, 1857: *Journal of the proceedings of the Linnean Society of*

London, *Zoology* 2: 76, (w.) Type location: Singapore.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Kao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Trang*: Khao Chong.

***Crematogaster inflata* F. Smith, 1857**

มดฮี้โป่งเหลือง

Crematogaster inflata F. Smith, 1857: *Journal of the Proceedings of Linnean Society of London, Zoology* 2: 76, pl. 2 fig. 2 (w.) Type locality: Singapore.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui; *Phrae*: Wang Chin; *Phitsanulok*; Tung Salaeng Luang.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Sakaerat.

CENTRAL: *Saraburi*: Phu Kae.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chantaburi*: Nam Tok Phliu.

***Crematogaster modiglianii* Emery, 1900**

มดฮี้ปีกงัด

Crematogaster modiglianii Emery, 1900: *Annali del Museo Civico di Storia Naturale di Genova* (2) 20: 688 (w.) Type locality: Indonesia.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong; *Songkhla*: Ton Nga Chang, Khao Nam Khang; *Narathiwat*: Halabala, Toa Dang.

***Crematogaster rogenhoferi* Mayr, 1879**

มดฮี้ทุ่ง

Crematogaster rogenhoferi Mayr, 1879: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 28: 983 (w.) Type locality: Myrmmar.

DISTRIBUTION:

NORTHERN: *Phayao*: Mae Yluar; *Phrae*: Wang Chin; *Tak*: Thung Yai. *Phitsanulok*: Thung Salaeng Luang.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat; *Surin*: Haui Tab Tan.

CENTRAL: *Saraburi*: Phu Kae; *Bangkok*: Kasetsart University.

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Narathiwat*: Halabala, Toa Dang.

***DACETINOPS* (Myrmicinae: Stenammini)**

สกุลมดลี้ม

Dacatinops Brown and Wilson, 1957a: 1.

Type species: *Dacatinops cibdelus* Brown and Wilson, 1957, by original designation Revision of genus: Tayler, 1985: 41.

***Dacatinops concinnus* Tayler, 1965**

มดลี้ม

Dacatinops concinnus Tayler, 1965:

Brevora 237, figs. 1-2. (w.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Songkhla*: Khao Nam Khang.

***DIACAMMA* (Ponerinae: Ponerini)**

สกุลมดหนามคู้

Diacamma Mayr, 1862: 718. Type species:

Ponera rugosa Le Guillou, 1842, by subsequent designation of Bingham, 1903:75.

***Diacamma intricatum* (F. Smith, 1857)**

มดหนามคู้บอร์เนียว

Ponera intricata F. Smith, 1857: *Journal*

of the proceedings of the Linnean Society of London, Zoology 2: 67 (w.) Type locality: Borneo.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Nam Tok Phliu.

***Diacamma rugosum* (Le Guillou, 1842)**

มดหนามคู้สี่เทา

Ponera rugosa Le Guillou, 1842: *Annales*

de la Societe Entomologique de France 10: 318 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Lampang*: Nago.

NORTHEASTERN: *Kalasin*: Phu Si Than.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHWESTERN: *Uthai Thani*: Huai Kha Khang.

SOUTEASTERN: *Sa Kaeo*: Nam Tok Khao Takrub; *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Narathiwat*: Halabala; *Trang*:
Khao Chong.

***Diacamma sculpturata* (F. Smith, 1859)**

มดหนามคู้ท้องร้อง

Ponera sculpturata F. Smith, 1859: *Journal of the proceedings of the Linnean Society of London, Zoology* 3: 142 (w.) Type locality: Indonesia.

DISTRIBUTION:

NORTHERN: *Lampang*: Mae Had; *Phrea*: Wang
Chin.

PENINSULAR: *Trang*: Khao Chong; *Songkhla*:
Khao Nam Khang; *Narathiwat*:
Halabala.

***Diacamma vargens* (F. Smith, 1860b)**

มดหนามคู้ดำ

Ponera vargens F. Smith, 1860b: *Journal of the proceedings of the Linnean Society of London, Zoology* 5 (supplement to Volume4): 103 (w.) Type locality: Indonesia.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao, Mae
Cham, Mae Tang; *Phrae*: Wang
Chin.

EASTERN: *Chaiyaphum*: Phu Khieo, *Nakhon
Ratchasima*: Khao Yai, Sakaerat.

CENTRAL: *Saraburi*: Phu Kae; Bangkok:
Kasetsart University.

SOUTHEASTERN: *Chachoensao*: Khao Ang
Rei Nai.

PENINSULAR: *Narathiwat*: Halabala.

DOLICHODERUS (Dolichoderinae)

สกุลมดกันท้อย

Dolichoderus Lund, 1831: 130. Type species: *Formica attelaboides* Fabricius, 1775, by monotypy. *Dolichoderus* senior synonym of *Hypoclinea* and *Monacis*: Forel, 1878: 386. *Dolichoderus* senior synonym of *Acanthoclinea* and *Diceratoclinea*: Taylor and Brown, D. R. 1985: 93. *Dolichoderus*: senior synonym of *Acanthoclinea*, *Diabolus*, (junior homonym), *Diceratoclinea*, *Hypoclinea*, *Karawajewella*, *Monacis* and *Monoceratoclinea*, and Review of genus: Shattuck, 1992b: 66. Key to Australian species: Clark, 1930: 252. Key to Neotropical species: Kempf, 1959: 227;

Kempf, 1969: 292; Kempf, 1972: 253; Lattke, 1987: 263; Mackey, 1993; Shattuck, 1994: 39.

***Dolichoderus becarii* Emery, 1887a.**

มดกันท้อยท้ายแหลม

Dolichoderus becarii Emery, 1887a: *Annali del Museo Civico di Storia Naturale de Genova* (2) 4: 253 (q.) Type locality: Sumatra.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

***Dolichoderus cuspidatus* (F. Smith, 1857)**

มดกันท้อยอกหนาม

Polyrhachis cuspidatus F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 62 (w.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong; *Narathiwat*:
Halabala, Toa Dang.

***Dolichoderus semirugosus* (Mayr, 1870)**

มดกันท้อยยอดไม้

Hypoclinea semirugosus Mayr, 1870: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 20: 956 (diagnosis in key) (w.) Type locality: Borneo.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai.

***Dolichoderus sulcaticeps* (Mayr, 1870)**

มดกันท้อยเอวกลม

Hypoclinea sulcaticeps Mayr, 1870: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 20: 957 (diagnosis in key) (w.) Type locality: Borneo.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoensao*: Khao Ang
Reu Nai.

PENINSULAR: *Songkhla*: Hat Yai.

***Dolichoderus thoracicus* (F. Smith, 1860b)**

มดกันท้อยธรรมดา

Tapinoma thoracicus F. Smith, 1860b: *Journal of the proceedings of the Linnean Society of London, Zoology* 5: 69 (w.) Type Locality: Sulawesi.

DISTRIBUTION:

Found in all provinces.

***Dolichoderus tuberifer* Emery, 1887a**

มดกันท้อยหลังง่าม

Dolichoderus tuberifer Emery, 1887a: *Annali del Museo Civico di Storia Naturale de Genova* (2) 4: 253 (q.) Type locality: Sumatra.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao, Doi Ang Khang, Doi Suthep-Pui, Doi Inthanon, Mae Cham, Mae Tang, Doi Luang, Ob Luang; *Chiang Rai*: Mae Pha Luang; *Phayao*: Mae Ka; *Lampang*: Tham Pha Thai, Ngao; *Phrae*: Wang Chin; *Tak*: Thung Yai, Lan Sang; *Phitsanulok*: Thung Salaeng Luang.

NORTHEASTERN: *Phetchabun*: Khao Khor; *Udonthani*: Nong Sang; *Kalasin*: Phu Si Than; *Khon Khaen*: Chum Sang.

EASTERN: *Chaiyaphum*: Phu Khieo; Bamnead Narong, *Nakhon Ratchasima*: Khao Yai, Sakaerat; *Surin*: Huai Tab Tan; *Ubon Ratchathani*: Khong Chiam.

SOUTHWESTERN: *Uthai Thani*: Huai Kha Khang; *Kanchanaburi*: Thong Pha Phum, Salak Pra, Sai Yok; *Prachuap Khiri Khan*: Huai Yang.

CENTRAL: *Saraburi*: Phu Kea.

SOUTHEASTERN: *Sa Kaeo*: Nam Tok Khao Takrub; *Chachoengsao*: Khao Ang Reu Nai; *Chon Buri*: Khao Ang Reu Nai; *Rayong*: Khao Ang Reu Nai; *Chanthaburi*: Khao Ang Reu Nai, Khao Soi Dao, Nam Tok Phliu; *Trat*: Ko Chang.

PENINSULAR: *Chumphon*: Krom Luang Chumphon; *Ranong*: Klong Yan, Klong Naka; *Surat Thani*: Klong Sang; *Phangnga*: Khao Lak; *Nakhon Si Thammarat*: Khao Luang; *Phatthalung*: Thale Noi; *Trang*: Khao Chong; *Satun*: Thale Ban; *Songkhla*: Khao Nam Khang, Khao Khor Hong, Ton Nga

Chang; *Narathiwat*: Halabala, Toa Dang.

***DORYLUS* (Dorylinae)**

สกุลเสี้ยนดิน

Dorylus Fabricius, 1793: 365. Type species: *Vespa helvola* Linnaeus, 1764, by monotypy. Subgenera of *Dorylus* (alphabetical order with synonyms indicated): nominal plus *Alaopone*, *Anomma*, *Cosmaecetes*, (junior synonym of *Typhlopone*), *Dichthadia*, *Rhogmus*, *Shuckadia* junior synonym of *Alaopone*, *Sphegomyrmex* (junior synonym of *Anomma*), *Typhlopone*. Key to Indo-Australian species: Wilson, 1964; Gotwald, 1982: 157.

***Dorylus laevigatus* (F. Smith, 1857)**

เสี้ยนดินทุ่ง

Typhlopone laevigata F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 70 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Doi Chiang Dao.

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat.

PENINSULAR: *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala.

***Dorylus orientalis* Westwood, 1835**

เสี้ยนดินถั่ว

Dorylus orientalis Westwood, 1835: *Proceedings of the Zoology Society of London* 3: 72 (w.) Type locality: India.

DISTRIBUTION:

EASTERN: *Roi Et*.

***Dorylus vishnui* Wheeler, 1913b**

เสี้ยนดินป่า

Dorylus (Alaopone) vishnui Wheeler, 1913b: *Records of the Indian Museum* 8: 233 (w.) Type locality: Myanmar.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Rei Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Narathiwat*: Halabala.

ECHINOPLA (Formicinae: Camponotini)**สกุลมดมูม**

Echinopla F. Smith, 1857: 79. Type species: *Echinopla melanarctos* F. Smith, 1857, by subsequent designation of Wheeler, W. M. 1911: 162. *Echinopla* senior synonym of *Mesoxena*: Bolton, 1994: 50.

Echinopla melanactos* F. Smith, 1857*มดมูมขาดำ**

Echinopla melanactos F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 79, pl. 1, figs. 25-29 (w.) Type locality: Singapore.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

Echinopla pallipes* F. Smith, 1857*มดมูมขาเหลือง**

Echinopla pallipes F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 80 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

PENINSULAR: *Narathiwat*: Halabala.

EMERYOPONE (Ponerinae: Ponerini)**สกุลมดเขี้ยวเล็กน้อย**

Emeryopone Forel, 1912b: 761. Type species: *Emeryopone buttelreepeni* Forel, 1912b, by monotypy. *Emeryopone* junior synonym of *Belonopelta*: U. Baroni, 1975b: 296.

Emeryopone buttelreepeni* Forel, 1912b*มดเขี้ยวเล็กน้อย**

Emeryopone buttelreepeni Forel, 1912b: *Revue Suisse de Zoologie* 20: 762 (w.) Type locality: Sumatra.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Ranong*: Klong Naka; *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala.

EUPRENOLEPIS (Formicinae: Lasiini)**สกุลมดดอกคอด**

Euprenolepis Emery, 1906: 134. Type species: *Prenolepis procera* Emery, 1897, by original designation. *Euprenolepis* raised to genus and senior synonym of *Chapmanella*: Brown, 1953b: 6.

Euprenolepis procera* (Emery, 1897)*มดดอกคอดตาโต**

Rhopalothrix procera Emery, 1897: *Termeszetrajzi Fuzetek* 20: 572, pl.14, fig. 18 (w.q.) Type locality: New Guinea.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong; *Narathiwat*: Halabala.

GNAMPTOGENYS (Ponerinae: Ectatommini)**สกุลมดค่อม**

Gnamptogenys Roger, 1863a: 174. Type species: *Ponera tornata* Roger, 1861, subsequent designation of Emery, 1911b: 44. *Gnamptogenys* subgenus of *Ectatomma*: Mayr, 1887: 540; maintained as subgenus; Dalla Torre, 1893: 23; Emery, 1911b: 44; Wheeler, W.M. 1922: 644. *Gnamptogenys* revived status as genus, senior synonym of *Alfaria* (and its junior synonym of *Opisthoscyphus*), *Barbourella*, *Commateta*, *Emeryella*, *Holcoponera*, *Parectatomma*, *Poneracantha*, *Rhopalopone* (and its junior synonym of *Mictoponera*), *Spaniopone*, *Stictoponera*, *Tammotoca*, *Wheeleripone*, and revision of genus: Brown, 1958: 211. Key to New World species: Brown, 1958: 237.

Gnamptogenys bicolor* (Emery, 1889)*มดค่อมอกแดง**

Ectatomma (Stictoponera) bicolor Emery, 1889: *Annali del Museo Civico di Storia Naturale de Genova* (2) 7: 493 (w.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao, Doi Suthep- Pui, Doi Luang; *Tak*: Thung Yai; *Phrae*: Wang Chin.

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat; *Chaiyaphum*: Phu Khieo.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Trang*: Khao Chong, Palian;
Narathiwat: Halabala, Toa Dang.

***Gnamptogenys binghamii* (Forel, 1900)**

มดค่อมบึงแสม

Ectatomma (Stictoponera) binghamii
Forel, 1900: *Mittheilungen der Schweizerischen
Entomologischen Gesellschaft* 10: 317 (w.) Type
locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang;
Tak: Thung yai;

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon
Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu
Nai; *Chanthaburi*: Khao Soi
Dao, Nam Tok Phliu.

***Gnamptogenys costata* (Emery, 1889)**

มดค่อมห้องลาย

Ectatomma (Stictoponera) costatum
Emery, 1889: *Annali del Museo Civico di Storia
Naturale de Genova* (2) 7: 494 (w.) Type locality:
Myanmar.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang
Reu Nai.

***Gnamptogenys manadensis* (Mayr, 1887)**

มดค่อมป่า

Ectatomma (Stictoponera) manadensis
Mayr, 1887: *Verhandlungen der k.k. Zoologisch-
Botanischen Gesellschaft in Wien* 37: 539 (foot note)
(w.) Type locality: Sulawesi.

DISTRIBUTION.

PENINSULAR: *Ranong*: Klong Naka; *Narathiwat*:
Halabala.

HARPEGNATHOS (Ponerinae: Ponerini)

สกุลมดเขี้ยวโจ่ง

Harpegnathos Jerdon, 1851: 116. Type
species: *Harpegnathos saltator* Jerdon, 1851, by
monotypy. *Harpegnathos* senior synonym of
Drepanognathus: Roger, 1863b: 22. *Harpegnathos*
incorrectly as junior synonym of *Drepanognathus*:
Dalla Torre, 1893: 22. *Harpegnathos* senior synonym
of *Drepanognathus*: Forel, 1893a, 166; Forel, 1900b:
63.

***Harpegnathos venator* (F. Smith, 1858)**

มดเขี้ยวโจ่ง

Drepanognathus venator F. Smith, 1858:
*Catalogue of Hymenopterous Insects in the Collec-
tion of British Museum* 6 Formicidae: 82 (w.) Type
locality: India.

DISTRIBUTION:

NORTHERN: *Phitsanulok*: Thung Salaeng Luang.
EASTERN: *Nakhon Ratchasima*: Sakaerat.

IRIDOMYRMEX (Dolichoderinae)

สกุลมดดำ

Iridomyrmex Mayr, 1862: 702. Type spe-
cies: *Formica detecta* F. Smith, 1858 (junior synonym
of *Iridomyrmex purpureus*), by subsequent designa-
tion of Bingham, 1903: 279. Review of genus:
Shattuck, 1992a: 14; Shattuck, 1992b: 101. Key to
species: Shattuck, 1993a: 113; Shattuck, 1993b: 1303;
Shattuck, 1994: 99.

***Iridomyrmex anceps* (Roger, 1863a)**

มดดำทุ่ง

Formica anceps Roger, 1863a: *Berliner
Entomologische Zeitschrift* 7: 164 (w.) Type locality:
West Malaysia.

DISTRIBUTION:

Found in all provinces.

LEPTANILLA (Leptanillinae: Leptanillini)

สกุลมดมดเอวย้อย

Leptanilla Emery, 1870: 196: Type species:
Leptanilla revelierii Emery, 1870, by monotypy.
Leptanilla senior synonym of *Leptomesites*: Baroni
Urbani, 1977: 433. Revision of genus: Baroni Urbani,
1977: 427.

***Leptanilla thai* Baroni, 1977**

มดเอวย้อยไทย

Leptanilla thai Baroni, 1977: *Entomologica
Basiliensia* 2: 454, figs. 21,23 (w.) Type locality: Thailand.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong.

LEPTOGENYS (Ponerinae: Ponerini)

สกุลมดเล็บหวี

Leptogenys Roger, 1861: 41. Type species:
Leptogenys falcigera Roger, 1861, by subsequent

designation of Bingham, 1903. *Leptogenys* senior synonym of *Microbolbos*: Wilson, 1955b: 136. *Leptogenys* senior synonym of *Machaerogenys* and *Lobopelta*: Bolton, 1975: 240. *Leptogenys* senior synonym of *Odontopelta*: Taylor and Brown, D.R. 1925: 32. *Leptogenys* senior synonym of *Prionogenys*: Taylor, 1988: 33. Key to Melanesian species: Wilson, 1958: 113. Caledonian species: Wilson, 1958: 136; Fijian species: Mann, 1921: 426.

***Leptogenys birmana* Forel, 1900c**

มดเล็บทวีปมา

Leptogenys (Lobopelta) birmana Forel, 1900c. *Journal of the Bombay Natural History Society* 13: 310 (w.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Inthanon, Mae Cham, Mae Tang, Doi Chiang Dao; *Lampang*: Tham Pha Thai; *Tak*: Thung Yai, Lan Sang; *Phitsanulok*: Thung Salaeng Luang.

NORTHEASTERN: *Phetchabun*: Khao Khor; *Udon Thani*: Nong Seang; *Kalasin*: Phusithan.

EASTERN: *Chaiyaphum*: Phu Khieo; *Surin*: Huai Tab Tan; *Nakhon Ratchasima*: Sakaerat, Khao Yai; *Ubon Ratchathani*: Khong Cheam.

SOUTHWESTERN: *Uthai Thani*: Huai Kha Khang; *Kanchanaburi*: Thong Pha Phum, Sai Yok; *Prachuap Khiri Khan*: Huai Yang.

CENTRAL: *Saraburi*: Phu Kae.

SOUTHEASTERN: *Sa Kaeo*: Nam Tok Khao Takrub; *Chachoengsao*: Khao Ang Reu Nai; *Chon Buri*: Khao Ang Reu Nai, Siracha; *Rayong*: Khao Ang Reu Nai; *Chanthaburi*: Khao Ang Reu Nai, Khao Soi Dao, Nam Tok Phliu; *Trat*: Ko Chang.

PENINSULAR: *Chumphon*: Krom Luang Chumphon; *Ranong*: Klong Yan, Klong Naka; *Nakhon Si Thammarat*: Khao Luang; *Trang*: Palian, Khao Chong; *Songkhla*: Khao Nam Khang, Khao

Khao Hong, Ton Nga Chang; *Narathiwat*: Halabala, Toa Dang.

***Leptogenys bornnensis* Wheeler, 1919**

มดเล็บทวีปอร์เนียว

Leptogenys (Lobopelta) bornnensis Wheeler, 1919: *Bulletin of the Museum of Comparative Zoology at Harvard College* 63: 59 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHWESTERN: *Uthai Thani*: Huai Kha Khang.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala.

***Leptogenys chalybaea* (Emery, 1887)**

มดเล็บทวีปเซียคราม

Lobopelta chalybaea Emery, 1887: *Annali del Museo Civico di Storia Naturale de Genova* (2) 5: 432 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Leptogenys diminuta* (F. Smith, 1857)**

มดเล็บทวีปกอด

Ponera diminuta F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 69 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Inthanon, Mae Cham, Mae Tang, Doi Chiang Dao; *Lampang*: Tham Pha Thai; *Tak*: Thung Yai, Lan Sang; *Phitsanulok*: Thung Salaeng Luang.

NORTHEASTERN: *Phetchabun*: Khao Khor; *Udon Thani*: Nong Seang; *Kalasin*: Phu Si Than.

EASTERN: *Chaiyaphum*: Phu Khieo; *Surin*: Huai Tab Tan; *Nakhon Ratchasima*: Sakaerat, Khao Yai; *Ubon Ratchathani*:

Khong Cheam.

SOUTHWESTERN: *Uthai Thani*: Huai Kha Khang;
Kanchanaburi: Thong Pha
Phum, Sai Yok; *Prachuap*
Khiri Khan: Huai Yang.

CENTRAL: *Saraburi*: Phu Kae.

SOUTHEASTERN: *Sa Kaeo*: Nam Tok Khao
Takrub; *Chachoengsao*: Khao
Ang Reu Nai; *Chon Buri*:
Khao Ang Reu Nai, Siracha;
Rayong: Khao Ang Reu Nai;
Chanthaburi: Khao Ang Reu
Nai, Khao Soi Dao, Nam Tok
Phliu; *Trat*: Ko Chang.

PENINSULAR: *Chumphon*: Krom Luang Chumphon;
Ranong: Klong Yan, Klong Naka;
Nakhon Si Thammarat: Khao
Luang; *Trang*: Palian, Khao Chong;
Songkhla: Khao Nam Khang, Khao
Khor Hong, Ton Nga Chang;
Narathiwat: Halabala, Toa Dang.

***Leptogenys hysteric* Forel, 1900c**

มดเล็บหัวเียวกลม

Leptogenys (Lobopelta) hysteric Forel,
1900c: *Journal of the Bombay Natural History*
Society 13: 311 (w.) Type locality: Sri Lanka.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai.
SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu
Nai; *Chanthaburi*: Khao Soi Dao.
PENINSULAR: *Ranong*: Klong Naka; *Trang*: Khao
Chong; *Songkhla*: Ton Nga Chang.

***Leptogenys iridescens* (F. Smith, 1857)**

มดเล็บหัวแดงรู้ง

Ponera iridescens F. Smith, 1857: *Journal*
of the proceedings of the Linnean Society of London,
Zoology 2: 66 (w.) Type locality: Borneo.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang
Reu Nai.

***Leptogenys kitteli* (Mayr, 1870)**

มดเล็บหัวอกร่อง

Lobopelta kitteli Mayr, 1870: *Verhandlungen*
der k.k. Zoologisch-Botanischen Gesellschaft in Wien
20: 966 (w.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Inthanon, Mae
Cham, Mae Tang, Doi Chiang Dao;
Lampang: Tham Pha Thai; *Tak*:
Thung Yai, Lan Sang; *Phitsanulok*:
Thung Salaeng Luang.

NORTHEASTERN: *Phetchabun*: Khao Khor; *Udon*
Thani: Nong Seang; *Kalasin*:
Phu Si Than.

EASTERN: *Chaiyaphum*: Phu Khieo; *Surin*:
Huai Tab Tan; *Nakhon Ratchasima*:
Sakaerat, Khao Yai; *Ubon*
Ratchathani: Khong Cheam.

SOUTHWESTERN: *Uthai Thani*: Huai Kha Khang;
Kanchanaburi: Thong Pha
Phum, Sai Yok; *Prachuap*
Khiri Khan: Huai Yang.

CENTRAL: *Saraburi*: Phu Kae.

SOUTHEASTERN: *Sa Kaeo*: Nam Tok Khao
Takrub; *Chachoengsao*: Khao
Ang Reu Nai; *Chon Buri*:
Khao Ang Reu Nai, Siracha;
Rayong: Khao Ang Reu Nai;
Chanthaburi: Khao Ang Reu
Nai, Khao Soi Dao, Nam Tok
Phliu; *Trat*: Ko Chang.

PENINSULAR: *Chumphon*: Krom Luang
Chumphon; *Ranong*: Klong Yan,
Klong Naka; *Nakhon Si Thammarat*:
Khao Luang; *Trang*: Palian, Khao
Chong; *Songkhla*: Khao Nam
Khang, Khao Khor Hong, Ton Nga
Chang; *Narathiwat*: Halabala,
Toa Dang.

***Leptogenys kraepelini* Forel, 1905**

มดเล็บหัวลิ่มเล็ก

Leptogenys (Lobopelta) kraepelini Forel,
1905: *Jahrbuch der hamburgischen Wissenschaftlichen*
Austalten [Mitteilungen aus dem Naturhistorischen
Museum] 22: 5 (w.) Type locality: Java.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai,
Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang
Reu Nai.

PENINSULAR: *Trang*: Khao Chong; *Narathiwat*: Halabala.

***Leptogenys mutabilis* (F. Smith, 1861)**

มดเล็บหัวหน้าเรียบ

Ponera mutabilis F. Smith, 1861: *Journal of the proceedings of the Linnean Society of London, Zoology* 6: 45 (w.) Type locality: Sulawesi.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Ranong*: Klong Naka; *Narathiwat*: Halabala.

***Leptogenys myops* (Emery, 1887b)**

มดเล็บหัวเหลี่ยมเล็ก

Lobopelta myops Emery, 1887b: *Annali del Museo Civico di Storia Naturale de Genova* (2) 5: 432 (w.) Type locality: Java.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Ranong*: Klong Naka; *Nakhon Si Thammarat*: Khao Luang; *Narathiwat*: Halabala.

LIOMYRMEX (Myrmicinae: Metaponini)

สกุลมดบอด

Liomyrmex Mayr, 1865: 23. Type species: *Myrmica caeca* F. Smith, 1860b. *Liomyrmex* senior synonym of *Laparomyrmex*: Dalla Torre, 1893: 63. *Liomyrmex* senior synonym of *Promyrma*: Forel, 1913: 26. Review of genus: Ettershank, 1966: 152.

***Liomyrmex gestroi* (Emery, 1887b)**

มดบอดถุ่ไน

Laparomyrmex gestroi Emery, 1887b: *Annali del Museo Civico di Storia Naturale de Genova* (2) 5: 461, pl.2, fig. 16 (q.) Type locality: New Guinea.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

LOPHOMYRMEX (Myrmicinae: Pheidolini)

สกุลมดดอกแหลม

Lophomyrmex Emery, 1892a: 114. Type

species: *Oecodoma quadrispinosa* Jerdon, 1851, Monotypy. Revision of genus: Ettershank, 1966: 132. Review of genus: Rigato, 1994: 47.

***Lophomyrmex bedoti* Emery, 1893b**

มดดอกแหลมไพโร

Lophomyrmex bedoti Emery, 1893b: *Revue Suisse de Zoologie* 1: 192, pl. 8, fig. 17 (w.q.) Type locality: Sumatra.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang.

SOUTHWESTERN: *Kanchanaburi*: Sai Yok.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Ranong*: Klong Naka; *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Songkhla*: Ton Nga Chang; *Narathiwat*: Halabala;

***Lophomyrmex birmanus* Emery, 1893b**

มดดอกแหลมพม่า

Lophomyrmex birmanus Emery, 1893b: *Revue Suisse de Zoologie* 1: 192 (diagnosis in key) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui, Mae Cham, Mae Tang, Doi Chiang Dao, Doi Luang; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

***Lophomyrmex lucidus* Menozzi, 1930**

มดดอกแหลมนิล

Lophomyrmex bedoti var. *lucida* Menozzi, 1930: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 79: 328 (w.) Type locality: West Malaysia.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham.

EASTERN: *Chaiyaphum*: Phu Khieo.

SOUTHWESTERN: *Kanchanaburi*: Sai Yok.

PENINSULAR: *Trang*: Khao Chong; *Songkhla*: Natawee, Khao Nam Khang.

***Lophomyrmex striatulus* Rigato, 1994**

มดดอกแหลมเมืองจันท

Lophomyrmex striatulus Rigato, 1994: *Systematic Entomology* 19: 56, fig. 16, 17 (w.) Type

locality: Thailand.

DISTRIBUTION:

SOUTHEASTERN: *Chanthaburi*: Nam Tok Phliu.

MERANOPLUS (Myrmicinae: Meranoplini)

สกุลมดโล้ท์

Meranoplus F. Smith, 1854: 224. Type species: *Cryptocerus bicolor* Guerin-Meneville, 1844, by subsequent designation of Bingham, 1903: 166. Revision of Afrotropical species: Bolton, 1981: 48.

***Meranoplus bicolor* (Guerin-Meneville, 1844)**

มดโล้ท์บ้าน

Cryptocerus bicolor Guerin-Meneville, 1844: Insectes (1829-1838) Paris: 425 (w.) Type locality: India.

DISTRIBUTION:

Found in all provinces.(open area).

***Meranoplus castaneus* F. Smith, 1857**

มดโล้ท์เหลือง

Meranoplus castaneus F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 81, pl. 2, fig. 7 (q.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong; *SongKhla*: Khao Khor Hong, Ton Ngachang; *Narathiwat*: Halabala, Toa Dang

***Meranoplus laeviventris* Emery, 1889**

มดโล้ท์เขาสูง

Meranoplus laeviventris Emery, 1889: *Annali del Museo Civico di Storia Naturale de Genova* (2) 7: 506, pl. 10, fig. 16 (w.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui, Doi Ang Khang; *Tak*: Thung Yai.
EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Sakaerat.

***Meranoplus mucronatus* F. Smith, 1857**

มดโล้ท์ปีกชนิด

Meranoplus mucronatus F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 82, pl. 2, fig. 6 (w.) Type locality: West Malaysia.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

MONOMORIUM (Myrmicinae: Solenopsidini)

สกุลมดละเอียด

Monomorium Mayr, 1855: 452. Type species: *Monomorium minutum* Mayr, 1855, by monotypy. *Monomorium* senior synonym of *Lampromyrmex*: Wheeler, W.M. 1915: 45. *Monomorium* senior synonym of *Paraphacota*: Santschi, 1927: 243. *Monomorium* senior synonym of *Epixenus*: Brown and Wilson, 1957b: 244. *Monomorium* senior synonyms of *Corynomyrmex*, *Epixenus*, *Epoecus*, *Equestrimessor*, *Holcomyrmex*, *Ireneidris*, *Isolcomyrmex*, *Lampromyrmex* (and its junior synonym of *Mitara*), *Paraphacota*, *Parholcolmyrmex*, *Trichomyrmex*, *Wheeleria* (homonym), *Wheeleriella*, *Xenhyboma*, and *Xeromyrmex* Ettershank, 1966: 82. *Monomorium* senior synonym of *Chelaner* (and its junior synonyms of *Notomyrmex*, *Protholcomyrmex*, and *Schizopelta*), and *Syllophopsis*: Bolton, 1987: 287.

***Monomorium chinense* Santschi, 1925**

มดละเอียดจีน

Monomorium minutum var. *chinensis* Santschi, 1925: *Buletin de la Societe Vaudoise des Sciences Naturelles* 56: 86 (w.) Type locality: China

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang, Doi Ang Khang.
EASRERN: *Nakhon Ratchasima*: Khao Yai.
SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Monomorium destructor* (Jerdon, 1851)**

มดละเอียดทองคำ

Atta destructor Jerdon, 1851: *Journal of Literature and Science* 17: 108 (w.) Type Locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Mae Tang, Ob Luang; *Lampang*: Tham Pha Thai.
EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat; Surin: Huai Tab Tan.
CENTRAL: *Saraburi*: Phu Kae; *Bangkok*: Kasetsart University, Chatuchak.
SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

Monomorium floricola* (Jerdon, 1851)*มดละเอียดหัวท้ายดำ**

Atta floricola Jerdon, 1851: *Journal of Literature and Science* 17: 107 (w.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao; *Tak*: Thung Yai; *Phrae*: Wang Chin.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Sakaerat.

CENTRAL: *Saraburi*: Phu Kae; *Bangkok*: Kasetsart University, Chatuchak.

SOUTHWESTERN: *Kanchanaburi*: Thong Pha Phum, Linthin.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chon Buri*: Siracha; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong, Palian; *Narathiwat*: Halabala, Toa Dang.

Monomorium latinode* Mayr, 1872*มดละเอียดเหลือง**

Monomorium latinode Mayr, 1872: *Annali del Museo Civico di Storia Naturale de Genova* 2: 152 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Lampang*: Tham Pha Thai.

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.

Monomorium pharaonis* (Linnaeus, 1758)*มดละเอียดบ้าน**

Formica pharaonis Linnaeus, 1758: Editio 10: 580 (w.) Type locality: Egypt.

DISTRIBUTION:

Found in all provinces.(open area)

Monomorium sechellense* Emery, 1894*มดละเอียดเซเชลล์**

Monomorium fossulatum subsp. *sechellense* Emery, 1894: *Annales de la Societe Emtomologique de France* 63: 69, fig. (w.q.) Type locality: Seychelle IS.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

Monomorium talpa* Emery, 1911a*มดละเอียดจุดตาเล็ก**

Monomorium talpa Emery, 1911a: *Lorentz. Zoologie*: 256 (w.) Type locality: New Guinea.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

MYOPOPONE* (Ponerinae: Amblyoponini)*สกุลมดหน้าโทน**

Myopopone Roger, 1861: 49. Type species: *Myopopone maculata* Roger, 1861 (junior synonym of *Myopopone castanea*), by subsequent designation of Bingham, 1903:33. Revision of genus: Brown, 1960: 170

Myopopone castanea* (F. Smith, 1860b)*มดหน้าโทนกลีสนิม**

Amblyopone castaneus F. Smith, 1860b: *Journal of the proceedings of the Linnean Society of London, Zoology* 5: 105, pl. 1, fig. 6 (w.) Type locality: Indonesia.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai; *Phitsanulok*: Thung Salaeng Luang.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao; *Trat*: Ko Chang.

PENINSULAR: *Trang*: Khao Chong; *Songkhla*: Ton Nga Chang.

MYRMICA* (Myrmicinae: Myrmicini)*สกุลมดเมืองหนาว**

Myrmica Latreille, 1804:179. Type species: *Formica rubra* Linnaeus, 1758: 580, by subsequent designation of Latreille, 1810: 437. *Myrmica* provisional senior synonyms of *Dodecamyrmica*, *Paramyrmica*, *Sifolinia*, *Sommimyrmica*, and *Symbiomyrmica*: Brown, 1973: 180-185. *Myrmica* senior synonym of *Dodecamyrmica*: Francoeur, 1981: 759.

Myrmica ritae* Emery, 1889*มดเมืองหนาวอินทนนท์**

Myrmica ritae Emery, 1889: *Annali del Museo Civico di Storia Naturale de Genova* (2) 7:

501, pl. 11, fig. 27 (w.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Inthanon.

MYRMICARIA (Myrmicinae: Myrmicariini)

สกุลมดหนวดเจ็ด

Myrmicaria Saunders, W.M. 1842: 57. Type species: *Myrmicaria brunnea* Saunders, W.M. 1842.

***Myrmicaria birmana* Forel, 1902**

มดหนวดเจ็ดยอดไม้

Myrmicaria arachnoides r. birmana Forel, 1902: *Revue Suisse de Zoologie* 10: 243 (w.) Type locality: Myanmar.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Myrmicaria brunnea* Saunders, W.M. 1842**

มดหนวดเจ็ดทุ่ง

Myrmicaria brunnea Saunders, W.M. 1842: *Transactions of the Entomological Society of London* 3: 57, pl. 5, fig.2 (m.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Ang Khang, Ob Luang, Doi Suthep-Pui, Mae Tang; *Phrae*: Wang Chin; *Tak*: Thung Yai; *Phitsanulok*: Thung Salaeng Luang.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Sakaerat, Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Ranong*: Klong Naka; *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala.

MYRMOTERAS (Myrmoteratini: Formicinae)

สกุลมดตาโปน

Myrmoteras Forel, 1893c: 607. Type species: *Myrmoteras binghamii* Forel, 1893c, by monotypy. Revisions of genus: Moffett, 1985: 1; Agosti, 1992: 405.

***Myrmoteras binghamii* Forel, 1893c**

มดตาโปนบึงแฮม

Myrmoteras binghamii Forel, 1893c: *Annales del la Societe Entomologique de Belgique* 37:

608 (w.) Type locality: Myanmar.

DISTRIBUTION:

Unknown area.(see Agosti, 1992: 409).

ODONTOMACHUS (Ponerinae: Ponerini)

สกุลมดกระโดด

Odontomachus Latreille, 1804: 174. Type species: *Formica haematoda* Linnaeus, 1758, by monotypy. *Odontomachus* senior synonym of *Pedetes*: Dalla Torre, 1893: 51. *Odontomachus* senior synonym of *Champsomyrmex* and *Myrtoterax*, and revision of genus: Brown, 1976: 67.

***Odontomachus latidens* Mayr, 1867**

มดกระโดดชวา

Odontomachus latidens Mayr, 1867: *Tijdschrift voor Entomologie* (2) 2: 80 (q.) Type locality: Java.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

***Odontomachus monticola* Emery, 1892b**

มดกระโดดเขาสูง

Odontomachus monticola Emery, 1892b: *Annales del la Societe Entomologique de France* 60: 560 (diagnosis in key) (w.) Type locality: Viet Nam.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Ang Khang, Ob Luang

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Satchasima*: Khao Yai.

***Odontomachus rixosus* F. Smith, 1857**

มดกระโดดเหลือง

Odontomachus rixosus F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 64 (w.) Type locality: Singapore.

DISTRIBUTION:

Found in all provinces.

***Odontomachus simillimus* F. Smith, 1858**

มดกระโดดบ้าน

Odontomachus simillimus F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 80, pl.5, fig. 8,9 (q.) Type locality: Fiji IS.

DISTRIBUTION:

Found in all provinces.

ODONTOPONERA (Ponerinae: Ponerini)**สกุลมดไ้ขึ้น**

Odontoponera Mayr, 1862: 717. Type species: *Ponera denticulata* (junior synonym of *Odontoponera transversa*), by monotypy.

Odontoponera denticulata* (F. Smith, 1858)*มดไ้ขึ้นดำ**

Ponera denticulata F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 90, pl.6, fig. 13, 14 (q.) Type locality: Singapore.

DISTRIBUTION:

Found in all provinces.

Odontoponera transversa* (F. Smith, 1857)*มดไ้ขึ้นมลายู**

Ponera transversa F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 68 (w.) Type locality: Singapore.

DISTRIBUTION:

PENINSULAR: *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala.

OECOPHYLLA (Oecophyllini: Formicinae)**สกุลมดแดง**

Oecophylla Smith, F. 1860b: 101. Type species: *Formica virescens* Fabricius, 1775 (junior synonym of *Oecophylla smaragdina*), by subsequent designation of Bingham, 1903: 310.

Oecophylla smaragdina* (Fabricius, 1775)*มดแดง หรือ มดแดงส้ม**

Formica smaragdina Fabricius, 1775: *Systema Entomologiae, sistens insectorum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, and observationibus*: 828 (q.) Type Locality: India.

DISTRIBUTION:

Found in all provinces.

PACHYCONDYLA (Ponerinae: Ponerini)**สกุลมดปุยฝ้าย**

Pachycondyla F. Smith, 1858: 105. Type species: *Formica crassinoda* Latreille, 1802, by subsequent designation of Emery, 1901a: 42.

Pachycondyla amblyops* (Emery, 1887b)*มดปุยฝ้ายเหลืองเล็ก**

Ponera amblyops Emery, 1887b: *Annali del Museo Civico di Storia Naturale de Genova* (2) 5: 434 (w.) Type locality: Sumatra.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham; *Tak*: Thung yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Narathiwat*: Halabala.

Pachycondyla astuta* F. Smith, 1858*มดปุยฝ้ายป่า**

Pachycondyla astuta F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 107 (w.) Type locality: Australia.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

Pachycondyla chinensis* (Emery, 1895)*มดปุยฝ้ายจ้วจิ้น**

Ponera nigrita subsp. *chinensis* Emery, 1895: *Annali del Museo Civico di Storia Naturale de Genova* (2) 14: 460 (w.) Type locality: China.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Luang; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHWESTERN: *Kanchanaburi*: Thong Pha Phum.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chon Buri*: Siracha; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Ranong*: Klong Naka; *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala.

Pachycondyla leeuwenhoekii* (Forel, 1886)*มดปุยฝ้ายออกร่อง**

Ponera leeuwenhoekii Forel, 1886: *Journal of the Asiatic Society of Bengal. Part 2. Natural Science* 55: 244 (w.) type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao, Mae Cham, Mae Tang; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*; *Khao Ang Reu Nai*; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Trang*: Khao Chong.

***Pachycondylla luteipes* (Mayr, 1862)**

มดปุยฝ้ายจั่วจุดตาสิบ

Ponera luteipes Mayr, 1862: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 12: 722 (w.q.) Type locality: Nicobar IS.

DISTRIBUTION:

Found in all provinces.

***Pachycondyla nigrita* (Emery, 1895)**

มดปุยฝ้ายจั่วใหญ่

Ponera nigrita Emery, 1895: *Annali del Museo Civico di Storia Naturale de Genova* (2) 14: 459 (w.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Luang, Doi Suthep-Pui, Mae Tang; *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat.

CENTRAL: *Bangkok*: Kasetsart University.

SOUTHEASTERN: *Chachoengsao*: *Khao Ang Reu Nai*; *Chanthaburi*: *Khao Soi Dao*, *Nam Tok Phliu*.

PENINSULAR: *Narathiwat*: *Toa Dang*, *Halabala*.

***Pachycondyla rubra* (F. Smith, 1857)**

มดปุยฝ้ายสีอิฐ

Ponera rubra F. Smith, 1857: *Journal of the Proceedings of the Linnean Society of London, Zoology* 2: 66 (q.) Type locality: Singapore.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: *Khao Ang Reu Nai*.

***Pachycondyla rufipes* (Jerdon, 1851)**

มดปุยฝ้ายขาตาลแดง

Ponera rufipes Jerdon, 1851: *Journal of Literature and Science* 17: 119 (w.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Lampang*: *Ngao*; *Phrae*: *Wang Chin*; *Tak*: *Thung Yai*.

SOUTHEASTERN: *Trat*: *Kho Chang*.

PENINSULAR: *Ranong*: *Klong Naka*; *Narathiwat*: *Halabala*.

PARATOPULA (Myrmicinae: Formicoxenini)

สกุลมดยอดไม้

Paratopula Wheeler, W.M. 1919: 144. Type species: *Atopomyrmex ceylonicus* Emery, 1901b, by original designation.

***Paratopula macta* Bolton, 1988**

มดยอดไม้

Paratopula macta Bolton, 1988: *Entomologists Monthly Magazine* 124: 140, fig. 2 (w.q.) Type locality: Borneo.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: *Khao Yai*.

CENTRAL: *Bangkok*: *Kasetsart University*.

SOUTHEASTERN: *Chachoengsao*: *Khao Ang Reu Nai*; *Chanthaburi*: *Khao Soi Dao*.

PARATRECHINA (Formicinae: Lasiini)

สกุลมดรำคาญ

Paratrechina Motschoulsky, 1863: 13. Type species: *Paratrechina currens* Motschoulsky, 1863 (junior synonym of *Paratrechina longicornis*), by subsequent designation of Wheeler, W.M. 1911: 170.

***Paratrechina longicornis* (Latreille, 1802)**

มดรำคาญขายาว

Formica longicornis Latreille, 1802: *Histoire Naturelle des Fourmis, et recueil de memoires et dobservations sur les abeilles, les araignees, les faucheurs insectes*: 113 (w.) Type locality: Senegal.

DISTRIBUTION:

Found in all provinces.

***Paratrechina opaca* (Emery, 1887a)**

มดรำคาญเหลือบฟ้า

Prenolepis clandestina var. *opaca* Emery, 1887a: *Annali del Museo Civico di Storia Naturale de Genova* (2) 4: 243 (w.) Type locality: Java.

DISTRIBUTION:

Found in all provinces.

PHEIDOLE (Myrmicinae: Pheidolini)**สกุลมดคัน**

Pheidole Westwood, 1839: 219. Type species: *Atta providens* Sykes, 1835, by monotypy.

Pheidole aglae* Forel, 1913*มดคันเอกลา**

Pheidole aglae Forel, 1913: *Zoologische Jahrbucher. Abteilung für Systematik, Geographie und Biologie der Tiere* 36: 32 (s.w.q.) Type locality: Java.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai.

PENINSULAR: *Narathiwat*: Halabala.

Pheidole annexus* Eguchi, 2001*มดคันบอร์เนียว**

Pheidole annexus Eguchi, 2001: *TROPICS Monograph Series* No. 2: 32, fig. 6 (w.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

Pheidole aristotelis* Forel, 1911a*มดคันไต้หวัน**

Pheidole aristotelis Forel, 1911a: *Revue Suisse de Zoologie* 19: 43 (s.w.m.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Ranong*: Klong Naka; *Trang*: Khao Chong; *Narathiwat*: Halabala.

Pheidole bluntschlii* Forel, 1911c*มดคันสุมาตรา**

Pheidole (Ceratopheidole) bluntschlii Forel, 1911c: *Bulletin de la Societe Vaudoise Sciences Naturelles* 47: 373 (w.) Type locality: Sumatra.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

Pheidole bugi* Wheeler, W.M. 1919*มดคันบ้าน**

Pheidole bugi Wheeler, W.M. 1919: *Bulletin of the Museum of Comparative Zoology at Harvard College* 63: 66 (s.w.) Type locality: Borneo.

DISTRIBUTION:

CENTRAL: *Bangkok*: Kasetsart University.

PENINSULAR: *Narathiwat*: Halabala.

Pheidole butteli* Forel, 1913*มดคันบัทเทิล**

Pheidole butteli Forel, 1913: *Zoologische Jahrbucher. Abteilung für Systematik, Geographie und Biologie der Tiere* 36: 36 (s.w.q.m.) Type locality: Sumatra.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong.

Pheidole capellinii* Emery, 1887b*มดคันหัวบวบ**

Pheidole capellinii Emery, 1887b: *Annali del Museo Civico di Storia Naturale de Genova* (2) 5: 463 (footnote), pl.1, fig. 14 (s.) Type locality: Java.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang, Mae Cham; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai.

Pheidole cariniceps* Eguchi, 2001*มดคันฐานแบน**

Pheidole cariniceps Eguchi, 2001: *TROPICS Monograph Series* No. 2: 41, fig. 10 (w.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

Pheidole clypeoconis* Eguchi, 2001*มดคันหน้าอ**

Pheidole clypeoconis Eguchi, 2001: *TROPICS Monograph Series* No. 2: 44, fig. 11 (w.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong; *Narathiwat*: Halabala.

Pheidole comata* F. Smith, 1858*มดคันใหญ่หัวขน**

Pheidole comata F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 176 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.

Pheidole dugasi* Forel, 1911d*มดคันใหญ่ท้องขอบขน**

Pheidole dugasi Forel, 1911d: *In Escherich,*

k. Termitenleben auf Ceylon: 222 (footnote) (w.) Type locality: Viet Nam.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Chiang Mai University.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

Pheidole elisae* Emery, 1900*มดคันหัวเรียบ**

Pheidole elisae Emery, 1900: *Annali del Museo Civico di Storia Naturale de Genova* (2) 20: 686 (s.) Type locality: Indonesia.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong.

Pheidole fervens* F. Smith, 1858*มดคันหัวกลม**

Pheidole fervens F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 176 (s.) Type locality: Singapore.

DISTRIBUTION:

CENTRAL: *Bangkok*: Kasetsart University.

SOUTHEASTERN: *Chachoengsao*: Khao ang Reu Nai.

PENINSULAR: *Trang*: Palian.

Pheidole gatesi* (Wheeler, W.M. 1927)*มดคันใหญ่เขาสูง**

Aphaenogaster (Atomyrma) gatesi Wheeler, W.M. 1927: *Psyche* 34: 44 (w.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui, Soi Ang Khang, Ob Luang; *Tak*: Thung Yai.

Pheidole hongkongensis* Wheeler, W.M. 1928a*มดคันฮ่องกง**

Pheidole rinae subsp. *hongkongensis* Wheeler, W.M. 1928a: *Bolletino del Laboratorio di Zoologia generale e agraria del R. Istituto Superiore agrario di Portici* 22: 11 (s.w.q.) Type locality: Hong Kong.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham.

EASTERN: *Nakhon Ratchasima*: Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

Pheidole hortensis* Forel, 1913*มดคันหน้ายักษ์**

Pheidole hortensis Forel, 1913: *Zoologische Jahrbucher. Abteilung fur Systematik, Geographie und Biologie der Tiere* 36: 38, fig.J (s.w.q.m.) Type locality: Java.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Sakaerat, Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Ranong*: Klong Naka; *Trang*: Khao Chong.

Pheidole incensa* Wheeler, W.M. 1928a*มดคันอินเซ็นส์**

Pheidole rinae subsp. *incensa* Wheeler, W.M. 1928a: *Bolletino del Laboratorio di Zoologia generale e agraria del R. Istituto Superiore agrario di Portici* 22: 13 (s.w.) Type locality: China.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang, Ob Luang.

EASTERN: *Nakhon Ratchasima*: Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

Pheidole inornata* Eguchi, 2001*มดคันกระโดด**

Pheidole inornata Eguchi, 2001: *TROPICS Monograph Series* No. 2: 66, fig.22 (w.) Type locality: Borneo.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Sakaerat, Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

SOUTHWESTERN: *Kanchanaburi*: Thong Pha Phum.

PENINSULAR: *Ranong*: Klong Naka; *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Natathiwat*: Halabala.

Pheidole inscrobiculata* Viehmeier, 1916*มดคันหนวดสั้น**

Pheidole inscrobiculata Viehmeier, 1916: *Archiv fur Naturgeschichte* 81: 120 (s.w.) Type locality: West Malaysia.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

Pheidole jacobsoni* Forel, 1911b*มดคันจำชวา**

Pheidole javana subsp. *jacobsoni* Forel, 1911b: *Notes from the Leyden Museum* 33: 203 (s.w.q.m.) Type locality: Java.

DISTRIBUTION:

SOUTHEASTERN: *Chanthaburi*: Nam Tok Phliu.

Pheidole longipes* (F. Smith, 1857)*มดคันชยาวาว**

Myrmica longipes F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 70, pl.1, fig. 6 (w.) Type locality: Singapore.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Songkhla*: Ton Nga Chang; *Narathiwat*: Halabala.

Pheidole magretti* Emery, 1887b*มดคันมาเกรทที**

Pheidole magretti Emery, 1887b: *Annali del Museo Civico di Storia Naturale de Genova* (2) 5: 461 (w.s.) Type locality: Java.

DISTRIBUTION:

PENINSULAR: *Ranong*: Klong Naka.

Pheidole megacephala* F. Smith, 1860*มดคันทุ่ง**

Pheidole megacephala F. Smith, 1860: *Journal of the Proceedings of the Linnean Society, Zoology* 5: 112 (w.) Type locality: Indonesia.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong.

Pheidole noda* F. Smith, 1874*มดคันอกร่อง**

Pheidole nodus F. Smith, 1874: *Transection of the Entomological Society of London* (4) 7: 407 (s.) Type locality: Japan.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang.

Pheidole nodifera* (F. Smith, 1858)*มดคันครีบบขน**

Atta nodifer F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 165 (w.) Type locality: China.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang, Doi Luang; *Tak*: Thung Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Narathiwat*: Halabala.

Pheidole parva* Mayr, 1865*มดคันแคระ**

Pheidole parva Mayr, 1865: *Zoologischer Theil. Formicidae*: 98, pl.4, fig.28 (s.w.) Type locality: Sri Lanka.

DISTRIBUTION:

CENTRAL: *Bangkok*: Kasetsart University.

SOUTHWESTERN: *Kanchanaburi*: Thong Pha Phum.

Pheidole pieli* Santschi, 1925*มดคันหัวแบน**

Pheidole pieli Santschi, 1925: *Bulletin de la Societe Vaudoise des Sciences Naturelles* 56: 83 (w.s.) Type locality: China.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

Pheidole plagiaria* F. Smith, 1860b*มดคันร่องโค้ง**

Pheidole plagiaria F. Smith, 1860b: *Journal of the Proceedings of the Linnean Society, Zoology* 5: 112 (s.w.) Type locality: Indonesia.

DISTRIBUTION:

Found in all provinces.

Pheidole planifrons* Santschi, 1920*มดคันหน้าแบน**

Pheidole planifrons Santschi, 1920: *Annales de la Societe Entomologique de Belgique* 60: 166, fig.1 (s.w.) Type locality: Viet Nam.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao, Mae Tang, Mae Cham; *Tak*: Thung Yai.

EASTERN: *Chiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Trang*: Khao Chong; *Narathiwat*: Halabala.

***Pheidole protea* Forel, 1912c**

มดคันโปรเตีย

Pheidole javana subsp. *protea* Forel, 1912c: *Zoologische Jahrbucher supplement* 15: 55 (s.) Type locality: Sumatra.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui.

***Pheidole quadricuspis* Emery, 1900**

มดคันออกหนาม

Pheidole quadricuspis Emery, 1900: *Annali del Museo Civico di Storia Naturale de Genova* (2) 20: 683 (w.s.) Type locality: Sumatra.

DISTRIBUTION:

PENINSULAR: *Songkhla*: Khao Nam Khang; *Narathiwat*: Halabala.

***Pheidole rabo* Forel, 1913**

มดคันราโบ

Pheidole rabo Forel, 1913: *Zoologische Jahrbucher. Abteilung fur Systematik, Geographie und Biologie der Tiere* 36: 28, fig. J (s.w.q.) Type locality: Sumatra.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Ob Luang, Mae Tang.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Pheidole rinae* Emery, 1900**

มดคันรีนา

Pheidole rinae Emery, 1900: *Annali del Museo Civico di Storia Naturale de Genova* (2) 20: 687 (s.) Type locality: Sumatra.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

PENINSULAR: *Trang*: Khao Chong.

***Pheidole sarawakana* Forel, 1911a**

มดคันซาราวัก

Pheidole sauberi subsp. *sarawakana* Forel, 1911a: *Revue Suisse de Zoologie* 19: 45 (s.w.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

***Pheidole sauberi* Forel, 1905**

มดคันหัวเลี่ยน

Pheidole sauberi Forel, 1905: *Jahrbuch der hamburgischen Wissenschaftlichen [Mitteilungen aus dem Naturhistorischen Museum]* 22: 18 (s.w.) Type locality: Java.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong; *Narathiwat*: Halabala.

***Pheidole smythiesii* Forel, 1902**

มดคันใหญ่ท้องเรียบ

Pheidole (Ceratopheidole) smythiesii Forel, 1902: *Revue Suisse de Zoologie* 10: 165 (s.) 185 (w.q.m.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham, Doi Suthep-Pui.

***Pheidole spathifera* Forel, 1902**

มดคันแผ่นห้อย

Pheidole spathifera Forel, 1902: *Revue Suisse de Zoologie* 10: 168 (s.) 187 (w.q.m.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Ang Khang, Mae Tang; *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Sakaerat.

***Pheidole taivanensis* Forel, 1912a**

มดคันไต้หวัน

Pheidole taivenensis Forel, 1912a: *Entomologische Mitteilungen* 1: 59 (s.) Type locality: Taiwan.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao.

***Pheidole tandjongensis* Forel, 1913**

มดคันทานโจ

Pheidole tandjongensis Forel, 1913: *Abteilung fur Systematik, Geographie und Biologie der Tiere* 36: 42 (s.w.) Type locality: Sumatra.

DISTRIBUTION:

Found in all provinces.

***Pheidole tjibodana* Forel, 1905**

มดคันจิบบดาส

Pheidole nodgii var. *tjibodana* Forel, 1905:

Jahrbuch der hamburgischen Wissenschaftlichen [Mitteilungen aus dem Naturhistorischen Museum]

22: 16 (s.w.q.) Type locality: Java.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai, Sakaerat.

***Pheidole tsailuni* Wheeler, 1929**

มดคันปากนกแก้ว

Pheidole tsailuni Wheeler, 1929: *Psyche* 36:

2. Type locality: Congo.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao yai, Sakaerat.

PENINSULAR: *Songkhla*: Khao Khor Hong; *Narathiwat*: Halabala.

***Pheidole yeensis* Forel, 1902**

มดคันจีน

Pheidole sulcaticeps r. *yeensis* Forel, 1902: *Revue Suisse de Zoologie* 10: 179 (s.), 187 (s.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao, Mae Tang, Mae Cham; *Tak*: Thung Yai.

EASTERN: *Chiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

***Pheidole zoceana* Santschi, 1925**

มดคันเมืองเหนือ

Pheidole nodgii var. *zoceana* Santschi, 1925: *Bulletin de la Societe Vaudoise des Sciences Naturelles* 56: 83 (w.s.) Type locality: China.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui.

PHEIDOLOGETON (Myrmicinae: Pheidologetonini)

สกุลมดง่าม

Pheidologeton Mayr, 1862: 750. Type species: *Oecodoma diversa* Jerdon, 1851, by subsequent designation of Bingham, 1903:160.

***Pheidologeton affinis* (Jerdon, 1851)**

มดง่ามจุดตาเล็ก

Oecodoma affinis Jerdon, 1851: *Madras Journal of Literature and Science* 17: 110 (s.w.) Type locality: India.

DISTRIBUTION:

Found in all provinces.

***Pheidologeton diversus* (Jerdon, 1851)**

มดง่ามทุ่ง

Oecodoma diversus Jerdon, 1851: *Madras Journal of Literature and Science* 17: 109 (s.w.) Type locality: India.

DISTRIBUTION:

Found in all provinces.

***Pheidologeton pygmaeus* Emery, 1887b**

มดง่ามดำ

Pheidologeton pygmaeus Emery, 1887b: *Annali del Museo Civico di Storia Naturale de Genova* (2) 5: 465 (w.s.) Type locality: Indonesia.

DISTRIBUTION:

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Rayong*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

***Pheidologeton silenus* (F. Smith, 1858)**

มดง่ามออกหนาม

Pheidole silenus F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 176 (s.w.) Type locality: Singapore.

DISTRIBUTION:

PENINSULAR: *Ranong*: Klongnaka; *Trang*: Khao Chong; *Songkhla*: Khao Nam Khang; *Narathiwat*: Halabala.

***Pheidologeton trechideros* Zhou and Zheng, 1997**

มดง่ามเมืองเหนือ

Pheidologeton trechideros Zhou and Zheng, 1997: *Zoological Research*, 18: 167, figs. 7-9 (w.) Type locality: China.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Ang Khang, Doi Inthanon.

PLATYTHYREA (Ponerinae: Platythyreini)**สกุลมดตลอด**

Platythyrea Roger, 1863a: 172. Type species: *Platythyrea punctata* F. Smith, 1858, by subsequent designation of Bingham, 1903: 73. *Platythyrea* senior synonym of *Eubothroponera* and revision of genus: Brown, 1975: 6.

Platythyrea parallela* (F. Smith, 1859)*มดตลอดท่อน**

Ponera parallela F. Smith, 1859: *Journal of the Proceedings of the Linnean Society, Zoology* 5: 143 (w.) Type locality: Indonesia.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Inthanon; *Lampang*: Ngao; *Tak*: Thung Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Narathiwat*: Halabala, Toa Dang.

Platythyrea quadridenta* Donisthorpe, 1941a*มดตลอดสี่ง่าม**

Platythyrea quadridenta Donisthorpe, 1941a: *Annals and Magazine of Natural History* (11) 7: 134 (w.) Type locality: New Guinea.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

Platythyrea tricuspadata* Emery, 1900*มดตลอดสามง่าม**

Platythyrea tricuspadata Emery, 1900: *Annali del Museo Civico di Storia Naturale de Genova* (2) 20: 665 (w.) Type Locality: Sumatra.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Halabala.

POLYRHACHIS (Formicinae: Camponotini)**สกุลมดหนาม**

Polyrhachis F. Smith, 1857: 58: Type species: *Formica bihamata* Drury, 1773, by original designation.

Polyrhachis abdominalis* F. Smith, 1858*มดหนามกระดิ่งแดง**

Polyrhachis abdominalis F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 63 (s.w.) Type locality: Myanmar. Subgenus: *Myrmhopla*.

DISTRIBUTION:

NORTHERN: *Phrae*: Wangchin; *Tak*: Thung Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Narathiwat*: Toa Dang, Halabala.

Polyrhachis arcuata* (Le Guillou, 1842)*มดหนามทึบเรียบ**

Formica arcuata Le Guillou, 1842: *Annales de la Societe Entomologique de France* 10: 313 (q.) Type locality: Philippines. Subgenus: *Charimyrmica*.

DISTRIBUTION:

CENTRAL: *Bangkok*: Bang Khen.

PENINSULAR: *Narathiwat*: Halabala.

Polyrhachis armata* (Le Guillou, 1842)*มดหนามกระดิ่งดำ**

Formica armata Le Guillou, 1842: *Annales de la Societe Entomologique de France* 10: 313 (q.) Type Locality: Philippines. Subgenus: *Myrmhopla*.

DISTRIBUTION:

Found in all provinces.

Polyrhachis bellicosa* F. Smith, 1859*มดหนามเคียวเล็ก**

Polyrhachis bellicosus F. Smith, 1859: *Journal of the Proceedings of the Linnean Society, Zoology* 5: 142 (w.) Type locality: Indonesia. Subgenus: *Polyrhachis*.

DISTRIBUTION:

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Ranong*: Klong Naka; *Narathiwat*: Halabala, Toa Dang.

Polyrhachis bicolor* F. Smith, 1858*มดหนามกระดิ่งไฟ**

Polyrhachis bicolor F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 65 (q.) Type locality: Myanmar. Subgenus: *Myrmhopla*.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang; *Phrae*: Wang Chin.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Narathiwat*: Halabala.

***Polyrhachis bihamata* (Drury, 1773)**

มดหนามเดี่ยวใหญ่

Formica bihamata Drury, 1773: *Illustrations of Natural History. Wherein are exhibited upwards of two hundred and twenty figures of exotic insects.* 3: 73, pl.38, figs.7, 8 (w.) Type locality: Madagascar. Subgenus: *Polyrhachis*.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao; *Lampang*: Tham Pha Thai; *Tak*: Thung Yai

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHWESTERN: *Uthai Thani*: Huai Kha Khang; *Kanchanaburi*: Thong Phaphum

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Songkhla*: Ton Nga Chang; *Narathiwat*: Halabala.

***Polyrhachis calypso* Forel, 1911c**

มดหนามกระตึงออกยาว

Polyrhachis spinosa subsp. *calypso* Forel, 1911c: *Bulletin de la Societe Vaudoise des Sciences Naturelles* 47: 394, (w.) Type locality: Sumatra. Subspecies: *Myrmhopla*.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Toa Dang.

***Polyrhachis dives* F. Smith, 1857**

มดหนามกระตึงขนทอง

Polyrhachis dives F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 64 (w.) Type locality: Singapore. Subgenus: *Myrmhopla*.

DISTRIBUTION:

Found in all provinces.

***Polyrhachis dolomedes* F. Smith, 1863**

มดหนามยอดไม้แดง

Polyrhachis dolomedes F. Smith, 1863: *Journal of the proceedings of the Linnean Society, Zoology* 6: 14 (w.q.) Type locality: Indonesia. Subgenus: *Myrmatopa*.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Polyrhachis flavicornis* F. Smith, 1857**

มดหนามยอดไม้ดำ

Polyrhachis flavicornis F. Smith, 1857: *Journal of the proceedings of the Linnean Society of London, Zoology* 2: 63 (w.) Type locality: Singapore. Subgenus: *Myrmatopa*.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Polyrhachis furcata* F. Smith, 1858**

มดหนามกระตึงเดี่ยว

Polyrhachis furcatus F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 64, pl. 4, fig. 20 (q.) Type locality: Myanmar. Subgenus: *Myrmhopla*.

DISTRIBUTION:

NORTHERN: *Phrae*: Wang Chin.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Nam Tok Phliu.

PENINSULAR: *Ranong*: Klong Naka; *Nakhon Si Thammarat*: Khao Luang; *Trang*: Khao Chong; *Narathiwat*: Halabala.

***Polyrhachi halidayi* Emery, 1889**

มดหนามปล้องใต้

Polyrhachi halidayi Emery, 1889: *Annali del Museo Civico di Storia Naturale de Genova* (2) 7: 517 (w.) Type locality: Myanmar. Subspecies: *Campomyrma*.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui, Doi Ang Khang, Mae Cham; *Phrae*: Wang Chin; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakaerat.

PENINSULAR: *Ranong*: Klong Naka; *Trang*: Khao Chong.

***Polyrhachis hauxwelli* Bingham, 1903**

มดหนามเกลี้ยงเฮวส์้อม

Polyrhachis hauxwelli Bingham, 1903: *Hymenoptera 2. Ants and Cuckowasps*: 394, fig. 133 (w.) Type locality: Myanmar.

DISTRIBUTION:PENINSULAR: *Trang*: Palian.***Polyrhachis hippomanes* F. Smith, 1861**

มดหนามอึบโปเล็ก

Polyrhachis hippomanes F. Smith, 1861: *Journal of the Proceedings of the Linnean Society, Zoology* 6: 43, pl. 1, fig. 21 (w.) Type locality: Sulawesi. Subgenus: *Myrmhopla*.

DISTRIBUTION:NORTHEASTERN: *Kalasin*: Phu Si Than.EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai, Sakhaerat.SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.PENINSULAR: *Songkhla*: Khao Nam Khang, *Narathiwat*: Toa Dang.***Polyrhachis illaudata* Walker, 1859**

มดหนามทึบทองแหลม

Polyrhachis illaudatus Walker, 1859: *Annals and Magazine of Natural History* (3) 4: 373 (w.) Type locality: Srilanka. Subgenus: *Myrma*.

DISTRIBUTION:

Found in all provinces.

***Polyrhachis inermis* F. Smith, 1858**

มดหนามทึบเอวขาว

Polyrhachis inermis F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 68, pl. 4, figs. 25, 26 (q.) Type locality: Sulawesi. Subgenus: *Myrma*.

DISTRIBUTION:SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.PENINSULAR: *Narathiwat*: Halabala.***Polyrhachis laevis* F. Smith, 1858**

มดหนามเกลี้ยงขาแดง

Polyrhachis laevis F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 64, pl. 4, figs. 42 (q.) Type locality: Myanmar. Subgenus: *Cyrtomyrma*.

DISTRIBUTION:

Found in all provinces.

***Polyrhachis latona* Wheeler, W.M. 1909**

มดหนามไต้หวัน

Polyrhachis latona Wheeler, W.M. 1909:

Bulletin of the American Museum of Natural History 26: 337 (w.) Type locality: Taiwan. Subgenus: *Myrma*.

DISTRIBUTION:EASTERN: *Nakhon Ratchasima*: Khao Yai.***Polyrhachis muelleri* Forel, 1893b**

มดหนามกระทิงเล็ก

Polyrhachis muelleri Forel, 1893b: *Journal of the Bombay Natural History Society* 8: 32 (w.) Type locality: Singapore. Subgenus: *Myrmhopla*.

DISTRIBUTION:SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.***Polyrhachis phalerata* Menozzi, 1926**

มดหนามอึบโปใหญ่

Polyrhachis (Myrmatopa) phalerata Menozzi, 1926: *Atti della Societa dei Naturalisti e Matematici di Modena* (6) 4: 102 (w.) Type locality: Philippines. Subgenus: *Myrmatopa*.

DISTRIBUTION:SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.***Polyrhachis proxima* Roger, 1863a**

มดหนามทึบทองง่าม

Polyrhachis proxima Roger, 1863a: *Berliner Entomologische Zeitschrift* 7: 155 (w.) Type locality: Indonesia. Subgenus: *Myrma*.

DISTRIBUTION:

Found in all provinces.

***Polyrhachis rastellata* (Latreille, 1802)**

มดหนามเกลี้ยงดำ

Formica rastellata Latreille, 1802: *Histoire Naturelle des Fourmis, et recueil de memoires et observations sur les abeilles, les araignees, les faucheurs insectes*: 130 (q.) Type locality: Indonesia. Subgenus: *Cyrtomyrma*.

DISTRIBUTION:NORTHERN: *Phrae*: Wang Chin; *Tak*: Thung Yai.PENINSULAR: *Ranong*: Klong Naka.***Polyrhachis risoxa* F. Smith, 1858**

มดหนามทึบดำขน

Polyrhachis risoxus F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 68 (q.) Type locality: Sulawesi. Subgenus: *Myrma*.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Polyrhachis striata* Mayr, 1862**

มดหนามทึบดำขอบไรขน

Polyrhachis striatus Mayr, 1862: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 12: 686, pl. 19, fig. 8 (w.) Type locality: Java. Subgenus: *Myrma*.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Polyrhachis sumatrensis* F. Smith, 1858**

มดหนามทึบดำสุมาตรา

Polyrhachis sumatrensis F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 65, pl. 4, fig 43 (q.) Type locality: Sumatra. Subgenus: *Myrma*.

DISTRIBUTION:

PENINSULAR: *Narathiwat*: Toa Dang, Halabala.

***Polyrhachis textor* F. Smith, 1857**

มดหนามตรีดำ

Polyrhachis textor F. Smith, 1857: *Journal of the Proceedings of the Linnean Society of London, Zoology* 2: 60 (w.) Type locality: West Malaysia. Subgenus: *Myrmothrinax*.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu nai.

***Polyrhachis thrinax* Roger, 1863a**

มดหนามตรีแดง

Polyrhachis thrinax Roger, 1863a: *Berliner Entomologische Zeitschrift* 7: 152 (w.) Type locality: Sri Lanka. Subgenus: *Myrmothrinax*.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

***Polyrhachis tibialis* F. Smith, 1858**

มดหนามกระตังเทา

Polyrhachis tibialis F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 63 (w.) Type locality: Myanmar. Subgenus: *Myrmhopla*.

DISTRIBUTION:

NORTHERN: *Phrae*: Wang Chin.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chon Buri*: Khao Nag Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Pattani*: Yaring; *Narathiwat*: Halabala.

***Polyrhachis ypsilon* Emery, 1887a**

มดหนามเขียวห่าง

Polyrhachis ypsilon Emery, 1887a: *Annali del Museo Civico di Storia Naturale de Genova* (2) 4: 239 (w.) Type locality: Singapore. Subgenus: *Polyrhachis*.

DISTRIBUTION:

PENINSULAR: *Ranong*: Klong Naka; *Narathiwat*: Halabala.

***PRENOLEPIS* (Formicinae: Lasiini)**

สกุลมดดอกคอด

Prenolepis Mayr, 1861: 52. Type species: *Tapinoma nitens* Mayr, 1853, by subsequent designation of Bingham, 1903: 325.

***Prenolepis jacobsoni* Crawley, 1923**

มดดอกคอดสุมาตรา

Prenolepis jacobsoni Crawley, 1923: *Entomologist's Record and Journal of Variation* 35: 30 (w.) Type locality: Sumatra.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Narathiwat*: Halabala.

***Prenolepis melanogaster* Emery, 1893b**

มดดอกคอดตาลท้องดำ

Prenolepis melanogaster Emery, 1893b: *Revue Suisse de Zoologie* 1: 223 (w.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Ang Khang, Mae Cham, Mae Tang.

***PRISTOMYRMEX* (Myrmicinae: Myrmecini)**

สกุลมดเต่า

Pristomyrmex Mayr, 1866b: 903. Type species: *Pristomyrmex pungens* Mayr, 1866b, by

monotypy.

***Pristomyrmex brevispinosus* Emery, 1887b**

มดเต่าทองแดง

Pristomyrmex brevispinosus Emery, 1887b: *Annali del Museo Civico di Storia Naturale de Genova* (2) 5: 451 (w.m.) Type locality: Sumatra.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui; *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

***Pristomyrmex pungens* Mayr, 1866b**

มดเต่าดำ

Pristomyrmex pungens Mayr, 1866b: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 16: 904, pl. 20, fig. 13 (w.) Type locality: West Malaysia.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Trang*: Khao Chong.

***Pristomyrmex trachylissus* F. Smith, 1858**

มดเต่าปีกชนิด

Pristomyrmex trachylissa F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of the British Museum* 6: 126 (q.) Type locality: Borneo.

DISTRIBUTION:

PENINSULAR: *Trang*: Khao Chong.

***PROATTA* (Myrmicinae: Stenammini)**

สกุลมดหลังหนาม

Proatta Forel, 1912b: 768. Type species: *Proatta butteli* Forel, 1912b, by monotypy.

***Proatta butteli* Forel, 1912b**

มดหลังหนาม

Proatta butteli Forel, 1912b: *Revue Suisse de Zoologie* 20: 769 (w.m.) Type locality: Sumatra.

DISTRIBUTION:

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon*

Ratchasima: Khao Yai, Sakaerat.

CENTRAL: *Saraburi*: Phu Kae; *Nakhon Nayok*: Khao Cha Ngok.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao, Nam Tok Phliu.

PENINSULAR: *Trang*: Khao Chong; *Narathiwat*: Halabala.

***PROBOLOMYRMEX* (Ponerinae: Platythyreini)**

สกุลมดไร้ตา

Probolomyrmex Mayr, 1901: 2. Type species: *Probolomyrmex filiformis* Mayr, 1901b, by monotypy.

***Probolomyrmex dammermani* Wheeler, W.M. 1928c**

มดไร้ตาชวา

Probolomyrmex dammermani Wheeler, W.M. 1928c: *Psyche* 35: 7, fig. 1 (w.) Type locality: Java.

DISTRIBUTION

NORTHERN: *Tak*: Thung Yai.

PENINSULAR: *Trang*: Khao Chong.

***PROCERATIUM* (Ponerinae: Proceratiini)**

สกุลมดค่อม

Proceratium Roger, 1863a: 171 Type species: *Proceratium silaceum* Roger, 1863a, monotypy.

***Procertium deelemani* Perrault, 1981**

มดค่อมอกหนาม

Procertium deelemani Perrault, 1981: *Nouvelle Revue d'Entomologie* 11: 189, figs. 1-6 (w.) Type locality: Borneo.

DISTRIBUTION

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Sa Keao*: Khao Ang Reu Nai.

***RECURVIDRIS* (Myrmicinae: Pheidologetonini)**

สกุลมดหนามกลับ

Recurvidris Bolton, 1992: 36. Replacement name for *Trigonogaster* Forel, 1890b: cviii.

***Recurvidris recurvispinosa* (Forel, 1890)**

มดหนามกลับ

Trigonogaster recurvispinosus Forel, 1890:

Annales de la Societe Entomologique de Belgique. Comptes-rendus 34: cix (w.) Type locality: India.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Sakaerat.
SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.
PENINSULAR: *Trang*: Khao Chong.

RHOPTROMYRMEX (Myrmicinae: Tetramoriini)

สกุลมดปลวก

Rhoptromyrmex Mayr, 1901: 18. Type species: *Rhoptromyrmex globinodis*, by subsequent designation of Wheeler, W.M. 1911: 172.

***Rhoptromyrmex wroughtonini* Forel, 1902**

มดปลวกทุ่ง

Rhoptromyrmex wroughtonini Forel, 1902: *Revue Suisse de Zoologie* 10: 231 (w.m.) Type locality: India.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Sakaerat, Khao Yai.
SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; Chanthaburi: Khao Soi Dao.

SOLENOPSIS (Myrmicinae: Solenopsidini)

สกุลมดคันไฟ

Solenopsis Wesrwood, 1840: 86. Type species: *Solenopsis mandibularis* Wesrwood, 1840, by monotypy.

***Solenopsis geminata* (Fabricius, 1804)**

มดคันไฟ

Atta geminata Fabricius, 1804: *Systema Piezatorum*: 423 (q.) Type Locality: Central America.

DISTRIBUTION:

Found in all provinces.(open area.)

TAPINOMA (Dolichoderinae)

สกุลมดเหม็น

Tapinoma Foerster, 1850: 43. Type species: *Tapinoma collina* Foerster, 1850, by monotypy.

***Tapinoma melanocephalum* (Fabricius, 1793)**

มดเหม็น

Formica melanocephala Fabricius, 1793: *Entomologia systematica emendata et aucta. Secun-*

dum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, and observationibus 2: 353 (w.) Type locality: French Guiana.

DISTRIBUTION:

Found in all provinces.

TECHNOMYRMEX (Dolichoderinae)

สกุลมดเอวแบน

Technomyrmex Mayr, 1872: 147. Type species: *Technomyrmex strenuus* Mayr, 1872, by monotypy.

***Technomyrmex albipes* (F. Smith, 1861)**

มดเอวแบนตีนขาว

Formica (Tapinoma) albipes F. Smith, 1861: *Journal of the proceedings of the Linnean Society, Zoology* 6: 38 (w.) Type locality: Sulawesi.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

***Technomyrmex butteli* Forel, 1913**

มดเอวแบนเล็ก

Technomyrmex butteli Forel, 1913: *Abteilung fur Systematik, Geographie und Biologie der Tiere* 36: 97, fig. C (m.w.q.) Type locality: West Malaysia.

DISTRIBUTION:

Found in all provinces.

***Technomyrmex kraepelini* Forel, 1905**

มดเอวแบนโคนขาขาว

Technomyrmex kraepelini Forel, 1905: *Jahrbuch der hamburgischen Wissenschaftlichen [Mitteilungen aus dem Naturhistorischen Museum]* 22: 23 (w.) Type locality: Java.

DISTRIBUTION:

Found in all provinces.

***Technomyrmex modiglianii* Emery, 1900**

มดเอวแบนหัวใจ

Technomyrmex modiglianii Emery, 1900: *Annali del Museo Civico di Storia Naturale de Genova* (2) 20 696, fig. 12 (w.) Type locality: Sumatra.

DISTRIBUTION:

Found in all provinces.

TETRAMORIUM (Myrmicinae: Tetramoriini)

สกุลมดรีว

Tetramorium Mayr, 1855: 423. Type spe-

cies: *Formica caespitum* Linnaeus, 1758, by subsequent designation of Girard, 1879: 1016.

***Tetramorium bicarinatum* (Nylander, 1846)**

มดรีวหน้าสัน

Myrmica bicarinata Nylander, 1846: *Acta Societatis Scientiarum Fennicae* 2: 1061 (w.q.) Type locality: U.S.A.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Ang Khang; *Phrae*: Wang Chin.

CENTRAL: *Bangkok*: Kasetsart University.

***Tetramorium ciliatum* Bolton, 1977**

มดรีวสยาม

Tetramorium ciliatum Bolton, 1977: *Bulletin of the British Museum (Natural History) (Entomology)* 36: 121, fig. 49 (w.q.) Type locality: Thailand.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Cham; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Sakaerat, Khao Yai.

***Tetramorium eleates* Forel, 1913**

มดรีวท้องดำ

Tetramorium (Xiphomyrmex) tortuosu var. *eleates* Forel, 1913: *Abteilung fur Systematik, Geographie und Biologie der Tiere* 36: 82 (w.q.) Type locality: Sumatra.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao.

EASTERN: *Nakhon Ratchasima*: Sakaerat, Khao Yai.

PENINSULAR: *Trang*: Khao Chong; *Songkhla*: Khao Nam Khang.

***Tetramorium flavipes* Emery, 1893c**

มดรีวขาเหลือง

Tetramorium (Xiphomyrmex) flavipes Emery, 1893c: *Annales de la Societe Entomologique de France* 62: 247 (footnote) (w.) Type locality: Thailand.

DISTRIBUTION:

NORTHERN: *Lampang*: Mae Had; *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Sakaerat.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi

Dao.

***Tetramorium insolens* (F. Smith, 1861)**

มดรีวสุลาเวสี

Myrmica insolens F. Smith, 1861: *Journal of the Proceedings of the Linnean Society, Zoology* 6: 47 (w.) Type locality: Sulawesi.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.

***Tetramorium kheperra* (Bolton, 1976)**

มดรีวทุ่ง

Triglyphothrix kheperra Bolton, 1976: *Bulletin of the British Museum (Natural History) (Entomology)* 34: 349, fig. 71 (w.) Type locality: Java.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang, Doi Suthep-Pui; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Sakaerat, Khao Yai.

PENINSULAR: *Songkhla*: Natawee; *Narathiwat*: Halabala.

***Tetramorium lanuginosum* Mayr, 1870**

มดรีวขาว

Tetramorium lanuginosum Mayr, 1870: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 20: 976 (w.) Type locality: Java.

DISTRIBUTION:

CENTRAL: *Bangkok*: Kasetsart University, Chatuchak.

EASTERN: *Nakhon Ratchasima*: Khao Yai.

SOUTHEASTERN: *Chanthaburi*: Panatnicom.

PENINSULAR: *Narathiwat*: Toa Dang.

***Tetramorium nipponense* Wheeler, W.M. 1928b**

มดรีวญี่ปุ่น

Tetramorium guineense subsp. *nipponense* Wheeler, W.M. 1928b: *Bollettino del Laboratorio di Zoologia generale e agraria del R. Istituto Superiore agrario di Portici* 22: 115 (w.) Type locality: Japan.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang; *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Sakaerat, Khao Yai.

Tetramorium palaense* Bolton, 1979*มดรีวอร์เนียว**

Tetramorium palaense Bolton, 1979: *Bulletin of the British Museum (Natural History) (Entomology)* 38: 175, fig. 56 (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang, Doi Chiang Dao; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Khao Yai.

Tetramorium parvispinum* (Emery, 1893b)*มดรีวหยาบขนปุย**

Trilyphothrix parvispina Emery, 1893b: *Revue Suisse de Zoologie* 1: 214 (diagnosis in key) (w.) Type locality: Borneo.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Chiang Dao.

Tetramorium parvum* Bolton, 1977*มดรีวท้องเหลือง**

Tetramorium parvum Bolton, 1977: *Bulletin of the British Museum (Natural History) (Entomology)* 36: 117 (w.) Type locality: Java.

DISTRIBUTION:

NORTHERN: *Tak*: Thung Yai.

EASTERN: *Nakhon Ratchasima*: Sakaerat, Khao Yai.

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.

Tetramorium simillimum* (F. Smith, 1851)*มดรีวเทศ**

Myrmica simillima F. Smith, 1851: *Hymenoptera Aculeata*: 118 (w.) Type locality: Great Britain.

DISTRIBUTION:

NORTHERN: *Lampang*: Tham Pha Thai.

EASTERN: *Nakhon Ratchasima*: Sakaerat, Khao Yai.

CENTRAL: *Bangkok*: Kasetsart University.

PENINSULAR: *Narathiwat*: Toa Dang.

Tetramorium smithi* Mayr, 1879*มดรีวสมิธ**

Tetramorium smithi Mayr, 1879: *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 28: 673 (w.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Mae Tang, Doi Inthanon.

EASTERN: *Nakhon Ratchasima*: Sakaerat, Khao Yai.

CENTRAL: *Bangkok*: Chatuchak.

PENINSULAR: *Narathiwat*: Halabala.

Tetramorium walshi* (Forel, 1890)*มดรีวเรียบขนปุย**

Trilyphothrix walshi Forel, 1890: *Annales de la Societe Entomologique de Belgique. Comptes-rendus* 34: cvii (w.q.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Ob Luang; *Phrae*: Wang Chin.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Sakaerat, Khao Yai.

TETRAPONERA (Pseudomyrmecinae)**สกุลมดตะนอย**

Tetraponera F. Smith, 1852: 44. Type species: *Tetraponera atrata* F. Smith, 1852 (junior synonym of *T. nigra*), by subsequent designation of Wheeler, W.M. 1911: 173.

Tetraponera allaborans* (Walker, 1859)*มดตะนอยคำเล็ก**

Pseudomyrma allaborans Walker, 1859: *Annals and Magazine of Natural History* (3) 4: 375 (w.q.m.) Type locality: Sri Lanka.

DISTRIBUTION:

NORTHERN: *Mae Hong Son*: Khun Yuam; *Chiang Mai*: Doi Suthep-Pui; *Chiang Rai*: Mae Rim; *Tak*: Thung Yai.

EASTERN: *Chaiyaphum*: Phu Khieo; *Nakhon Ratchasima*: Sakaerat, Khao Yai.

SUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai; *Chanthaburi*: Khao Soi Dao.

PENINSULAR: *Surat Thani*: Tai Rom Yen; *Phangnga*: Khao Lak; *Satun*: Thala Ban; *Trang*: Khao Chong.

Tetraponera attenuata* F. Smith, 1877*มดตะนอยคำใหญ่**

Tetraponera attenuata F. Smith, 1877: *Transactions of the Entomological Society of London* (4) 10: 71 (w.) Type locality: Borneo.

DISTRIBUTION:

Found in all provinces.

Tetraponera binghami* (Forel, 1902)*มดตะนอยบิงแฮม**

Sima binghami Forel, 1902: *Revue Suisse de Zoologie* 10: 243 (q.w.m.) Type locality: Myanmar.

DISTRIBUTION:

NORTHERN: *Mae Hong Son*: Khun Yuam; *Chiang Mai*: Doi Inthanon, Nam Tok Mae Ya.

Tetraponera crassiusculata* (Emery, 1900)*มดตะนอยหลังเกราะ**

Sima allaborans subsp. *crassiusculata* Emery, 1900: *Annali del Museo Civico di Storia Naturale de Genova* (2) 20: 677, fig. 6 (w.) Type locality: Sumatra.

DISTRIBUTION:

PENINSULAR: *Surat Thani*: Khao Sok.

Tetraponera difficilis* (Emery, 1900)*มดตะนอยดำขนทอง**

Sima difficilis Emery, 1900: *Annali del Museo Civico di Storia Naturale de Genova* (2) 20: 677, fig. 6 (w.) Type locality: Sumatra.

DISTRIBUTION:

SOUTHWESTERN: *Kanchanaburi*: Linthin
SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.
PENINSULAR: *Phangnga*: Khao Lak, Takua Pa; *Nakhon Si Thammarat*: Khao Laung, Nam Tok Phrom Lok; *Satun*: Thale Ban; *Songkhla*: Ton Nga Chang; *Yala*: Betong.

Tetraponera extenuata* Ward, 2001*มดตะนอยสิงคโปร์**

Tetraponera extenuata Ward, 2001: *Invertebrate Taxonomy* 15: 614, figs. 16, 33, 39 (w.) Type locality: Singapore.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Sakaerat.

Tetraponera modesta* (F. Smith, 1860b)*มดตะนอยเหลี่ยม**

Pseudomyrma modesta F. Smith, 1860b: *Journal of the Proceedings of the Linnean Society, Zoology* 5: 106 (w.) Type locality: Indonesia.

DISTRIBUTION:

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Satun*: Thale Ban.

Tetraponera nigra* (Jerdon, 1851)*มดตะนอยดำอินเดีย**

Eciton nigrum Jerdon, 1851: *Madras Journal of Literature and Science* 17: 112 (q.w.) Type locality: India.

DISTRIBUTION:

NORTHERN: *Chiang Mai*: Doi Suthep-Pui, Ob Luang; *Lampang*: Huai Tak.

PENINSULAR: *Satun*: Thale Ban.

Tetramorium nitida* (F. Smith, 1860b)*มดตะนอยท้องเรียบ**

Pseudomyrma nitida F. Smith, 1860b: *Journal of the Proceedings of the Linnean Society, Zoology* 5: 106 (w.) Type locality: Indonesia.

DISTRIBUTION:

SOUTHWESTERN: *Kanchanaburi*: Tham Than Lot.

EASTERN: *Si Sa Ket*: Kanthararom.

SOUTHEASTERN: *Chachoengsao*: Khao Ang Reu Nai.

PENINSULAR: *Phangnga*: Khao Lak; *Nakhon Si Thammarat*: Khao Luang; *Songkhla*: Hat Yai.

Tetraponera nodosa* Ward, 2001*มดตะนอยซาราวัก**

Tetraponera nodosa Ward, 2001: *Invertebrate Taxonomy* 15: 639, figs. 75, 83 (w.) Type locality: Sarawak.

DISTRIBUTION:

PENINSULAR: *Songkhla*: Hat Yai.

Tetraponera notabilis* Ward, 2001*มดตะนอยไทย**

Tetraponera notabilis Ward, 2001: *Invertebrate Taxonomy* 15: 640, figs. 76, 84 (w.) Type locality: Thailand.

DISTRIBUTION:

EASTERN: *Nakhon Ratchasima*: Sakaerat.

Tetraponera pilosa* (F. Smith, 1858)*มดตะนอยขนปุย**

Pseudomyrma pilosa F. Smith, 1858: *Catalogue of Hymenopterous Insects in the Collection of*

the British Museum 6: 160 (w.) Type locality: Borneo.

DISTRIBUTION:

SOUTHEASTERN: *Chanthaburi*: Khao Soi Dao.
PENINSULAR: *Phangnga*: Ton Takuapa;
Trang: Khao Chong; *Songkhla*:
Ton Nga Chang.

***Tetraponera rufonigra* (Jerdon, 1851)**

มดตะนอยอกส้ม

Eciton rufonigrum Jerdon, 1851: *Madras Journal of Literature and Science* 17: 111 (w.) Type locality: India.

DISTRIBUTION:

Found in all provinces.

ACKNOWLEDGMENTS

The authors would like to show their deep gratitude to the following: Professor Seiki Yamane, who kindly provided us ant specimens for comparison, and also supplied us with his list of ant collected in Thailand; Associate Professor Decha Wiwatwitaya of Kasetsart University for his constant and willing support to us and also be so kind to allow us to check the labelled specimens in the Ant Museum; Miss Sasitorn Hasin for her kindness of identification ant specimens; and lastly, Mr. Tadsana Jeentong for his industrious typing of the manuscript.

LITERATURE CITED

- Agosti, D. 1992. Revision of the ants genus *Myrmoteras* in the Malay Archipelago. *Revue Suisse de Zoologie* 99: 405-429.
- Baroni, U.C. 1973. Die Gattung *Xenometra*, ein objektives synonym. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 46: 199-201.
- Baroni, U.C. 1975a. Primi reperti del genere *Calyptomyrmex* Emery nel subcontinente Indiano. *Entomologica Basiliensia* 1: 395-411.
- Baroni, U.C. 1975b. Contributo alla conoscenza dei generi *Belonopelta* Mayr e *Leiopelta* gen.n. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 48: 293-310.
- Baroni, U.C. 1977. Materiali per una revisione della sottofamiglia *Leptanillinae* Emery. *Entomologica Basiliensia* 2: 427-488.
- Bernard, F. 1956a. Revision des *Leptothorax* d'Europe occidentale basées sur la biométrie et les genitales males. *Bulletin de la Societe Zoologique de France* 81: 151-165.
- Bernard, F. 1956b. Remarques sur le peuplement des Balears en fourmis. *Bulletin de la Societe d'Histoire Naturelle de l'Afrique du Nord* 47: 254-266.
- Bernard, F. 1956c. Revision des fourmis palearctiques du genre *Cardiocondyla* Emery. *Bulletin de la Societe d'Histoire Naturelle de l'Afrique du Nord* 47: 299-306.
- Bingham, C.T. 1903. *The fauna of British India, including Ceylon and Burma*. Hymenoptera 2. Ants and Cuckoo-wasps: 506 pp. London.
- Bolton, B. 1974. A revision of the Palaetropical arctic boreal ant genus *Cataulacus* Smith, F. *Bulletin of the British Museum (Natural History) (Entomology)* 30: 1-105.
- Bolton, B. 1975. A revision of the ant genus *Leptogenye* Roger in the Ethiopian region, with a review of the Malagasy species. *Bulletin of the British Museum (Natural History) (Entomology)* 31: 235-305.
- Bolton, B. 1976. The ant tribe Tetramoriini. Constituent genera, review of the smaller genera and revision of *Triglyphothrix* Forel. *Bulletin of the British Museum (Natural History) (Entomology)* 34: 281-379.
- Bolton, B. 1977. The ant tribe Tetramoriini. The genus *Tetramorium* Mayr in the Oriental and Indo-Australian region, and in Australia. *Bulletin of the British Museum (Natural History) (Entomology)* 36: 67-151.
- Bolton, B. 1979. The ant tribe Tetramoriini. The genus *Tetramorium* Mayr in Malagasy region and in the New World. *Bulletin of the British Museum (Natural History) (Entomology)* 38: 129-181.
- _____ 1981. A revision of the ant genera *Meranoplus* F. Smith, *Dicroaspis* Emery and *Calyptomyrmex* Emery in the Ethiopian zoogeographical region. *Bulletin of the British Museum (Natural History) (Entomology)* 42: 34-81.

- _____ 1982. Afrotropical species of the myrmicine ant genera *Cardiocondyla*, *Leptothorax*, *Melissotarsus*, *Essor* and *Cataulacus*. *Bulletin of the British Museum (Natural History) (Entomology)* 45: 307-370.
- _____ 1987. A revision of the *Solenopsis* genus-group and revision of Afrotropical *Monomorium* Mayr. *Bulletin of the British Museum (Natural History) (Entomology)* 54: 263-452.
- _____ 1988. A revision of *Paratopula* Wheeler, a forgotten genus of myrmicine ants. *Entomologists Monthly Magazine* 124: 125-143.
- _____ 1992. A review of the ant genus *Recurvidris*, a new name for *Trigonogaster* Forel. *Psyche* 99: 35-48.
- _____ 1994. *Identification Guide of the Ant Genera of the World*: 222 pp. Cambridge, Mass.
- _____ 1995. *A New General Catalogue of the Ants of the World*: 504 pp. Harvard University Press.
- _____ 1997. *Identification Guide to the Ant Genera of the World*. 222 pp. Harvard University Press.
- Brown, W.L., Jr. 1953a. Characters and synonymies among the genera of ants. Part 1. *Breviora* 11: 1-13.
- _____ 1953b. Characters and synonymies among the genera of ants. Part 2. *Breviora* 18: 1-8.
- _____ 1958. Contributions toward a reclassification of the Formicidae.2. Tribe Ectatommini. *Bulletin of the Museum of Comparative Zoology at Harvard College* 128: 175-362.
- _____ 1960. Contributions toward a reclassification of the Formicidae.3. Tribe Amblyoponini. *Bulletin of the Museum of Comparative Zoology at Harvard College* 122: 145-230.
- _____ 1963. Characters and synonymies among the genera of ants. Part 3. Some members of the tribe Ponerini. *Breviora* 190: 1-10.
- _____ 1973. A comparison of the Hylean and Congo-West African rain forest ant faunas. In Meggers, B.J., Ayensu, E.S. and Duckworth, W.D. (eds). *Tropical Forest Ecosystems in Africa and South America: a Comparative Review*: 161-185.
- _____ 1975. Contributions toward a reclassification of the Formicidae.5. Ponerinae, tribe Platythyreini, Cerapachyini, cylindromyrmecini, Acanthostichini, Aenictogitini. *Search Agriculture* 5. Entomology (Ithaca) 15: 1-115.
- _____ 1976. Contributions toward a reclassification of the Formicidae.6. Ponerinae, tribe Ponerini, subtribe Odontomachiti. Section A. Introduction, subtribal characters, genus *Odontomachus*. *Studia Entomologica* (N.S.) 19: 67-171.
- _____ 1978. Contributions toward a reclassification of the Formicidae.6. Ponerinae, tribe Ponerini, subtribe Odontomachiti. Section B. Genus *Anochetus* and bibliography. *Studia Entomologica* (N.S.) 20: 549-652.
- Brown, W.L., Jr. and E.O. Wilson, 1957a. *Dacetinops*, a new ant genus from New Guinea. *Breviora* 77: 1-7.
- _____ 1957b. A new parasitic ant of the genus *Monomorium* from Alabama, with a consideration of the status of genus *Epixenus* Emery. *Entomological News* 68: 239-246.
- Buren, W.F. 1959. A review of the species of *Crematogaster*, sensu stricto, in North America. Part 1. *Journal of the New York Entomological Society* 66: 119-134.
- Clark, J. 1930. The Australian ants of the genus *Dolichoderus*, subgenus *Hypoclinea* Mayr. *Australian Zoologist* 6: 252-268.
- Crawley, W.C. 1923. Myrmecological notes. *Entomologist's Record and Journal of Variation* 35: 29-32.
- Dalla Torre, C. G.de. 1893. *Catalogus Hymenopteroru, hucusque descriptorum systematicus et synonymicus* 7: 289 pp. Lipsias.
- Donisthorpe, H. 1941a. Description of new species of ant from New Guinea. *Annals and Magazine of Natural History* (11) 7: 129-144.
- _____ 1941b. Glanures myrmecologiques. *Entomologist's Record and Journal of Variation* 53: 36-38.
- _____ 1949. A species of *Calyptomyrmex* Emery from New Guinea. *Entomologists Monthly Magazine* 85: 186.
- Drury, D. 1773. *Illustrations of Natural History. Wherein are exhibited upwards of two hundred and twenty figures of exotic insects*

- 2: 90 pp. London.
- _____ 1782. *Illustrations of Natural History. Wherein are exhibited upwards of two hundred figures of exotic insects* 3: 76 pp. London.
- Eguchi, K. 2001. A Revision of the Bornean species of the Ant Genus *Pheidole* (Insecta: Hymenoptera: Formicidae: Myrmicidae). *TROPICS monograph series* No.2: 1-154.
- Emery, C. 1869. Descrizione di una nuova formica italiana. *Annuario del Museo Zoologico della R. Università di Napoli* 5: 117-118.
- _____ 1870. Studi mirmecologici. *Bullettino della Società Entomologica Italiana* 2: 193-201.
- _____ 1881. Spedizione Italiana nell' Africa Equatoriale. Risultati Zoologici. Formichi. *Annali del Museo Civico di Storia Naturale di Genova* (2) 16: 270-276.
- _____ 1884. Materiali per lo studio della fauna Tunisia raccolti da G. e L. Doria. 3 Rassegna della formiche della Tunisia. *Annali del Museo Civico di Storia Naturale di Genova* (2) 1: 373-386.
- _____ 1887a. Catalogo della formiche esistenti nelle collezioni del Museo Civico di Genova. Parte terza. Formiche della regione Indo-Malese e dell Australian. *Annali del Museo Civico di Storia Naturale di Genova* (2) 4: 209-258.
- _____ 1887b. Catalogo della formiche esistenti nelle collezioni del Museo Civico di Genova. Parte terza. Formiche della regione Indo-Malese e dell Australian (continuazione e fine). *Annali del Museo Civico di Storia Naturale di Genova* (2) 5: 427-473.
- _____ 1889. Viaggi di Leonardo Fea in Birmania e regioni vicini. 20. Formiche di Biernania e del Tenasserim raccolte da Leonardo Fea. *Annali del Museo Civico di Storia Naturale di Genova* (2) 7: 485-520.
- _____ 1890. Studi sulle formiche della fauna neotropica. *Bullettino della Società Entomologica Italiana* 22: 38-80.
- _____ 1892a. Sopra alcune formiche raccolte dall' Ingegnere L. Bricchetti Robecchi nel paese dei Somali. *Annali del Museo Civico di Storia Naturale di Genova* (2) 12: 110-122.
- _____ 1892b. Voyage de M. Ch. Alluaud dans le territoire d'Assinie (Afrique occidentale) en juillet et août 1886. Formicides. *Annales de la Société Entomologique de France* 60: 553-574.
- _____ 1893a. Untitled contribution introduced by, M.C. Emery, de Bologne, envoie les diagnoses de cinq nouveaux genres de Formicides. *Bulletin Bimensuel de la Société Entomologique de France* 1892 No. 20: cclxxv-cclxxvii.
- _____ 1893b. Voyage de MM. Bedot et Pictet dans l'Archipel Malais. Formicides de l'Archipel Malais. *Revue Suisse de Zoologie* 1: 187-229.
- _____ 1893c. Voyage de M.E. Simon a l'île de Ceylan (Janvier-février 1892). Formicides. *Annales de la Société Entomologique de France* 62: 239-258.
- _____ 1894. Mission Scientifique de M. Ch. Alluaud aux îles (mars, avril, mai 1892). Formicides. *Annales de la Société Entomologique de France* 63: 67-72.
- _____ C. 1895. Viaggio di Leonardo Fea in Birmania e regioni vicine. 63. Formiche di Birmania e dei Monti raccolte L.Fea. *Annali del Museo Civico di Storia Naturale di Genova* (2) 14: 450-483.
- _____ 1896. Formicides recoltés a Buitenzorg (Java), par M. Massart. *Annales de la Société Entomologique de France* 40: 245-249.
- _____ 1897. Formicidarum species novae vel minus cognitae collectione Musaei Nationalis Hungarici, quas in Nova Guinea, colonia germanica, collegit L. Biro. *Termesztudományi Füzetek* 20: 571-599.
- _____ 1900. Formiche raccolte da Elio Modigliani in Sumatra, Engano e Mentawai. *Annali del Museo Civico di Storia Naturale di Genova* (2) 20: 661-722.
- _____ 1901a. Note sur les sous-familles des dorylines et ponerines (famille des formicides). *Annales de la Société Entomologique de France* 45: 32-54.
- _____ 1901b. Ameisen gesammelt in Ceylon von Dr. W. Horn, 1899. *Deutsche Entomologische Zeitschrift* 1901: 113-122.
- _____ 1902. Note mirmecologiche. Rendiconto

- delle sessioni deela R. *Accademia delle Scienze dell'Istituto di Bologna (N.S.)* 6: 22-34.
- _____ 1906. Note sur *Prenolepis vividula* Nyl. Et sur la classification des especes du genre *Prenolepis*. *Annales de la Societe Entomologique de Belgique* 50: 130-134.
- _____ 1911a. Nova Guinea. Resultats de l'expedition scientifique neerlandaise a la Nouvelle-Guinee en 1907 et 1909 sous les auspices de Dr. H. A. Lorentz. 9 *Zoologie*: 249-259.
- _____ 1911b. In Wytsman, P. *Geneera Insectorum. Hymenoptera, Fam. Formicidae, subfam. Ponerinae. Fasc. 118*: 124 pp. Bruxelles.
- _____ 1912. Les especes-type des genres et sous-genres de la famille des formicides. *Annales de la Societe Entomologique de Belgique* 56: 271-273.
- _____ 1922. In Wytsman, P. *Geneera Insectorum. Hymenoptera, Fam. Formicidae, subfam. Myrmicinae. Fasc. 174B*: 95-206. Bruxelles.
- _____ 1925. In Wytsman, P. *Geneera Insectorum. Hymenoptera, Fam. Formicidae, subfam. Formicinae. Fasc. 183*: 302 pp. Bruxelles.
- Emery, C. and A. Foral. 1879. Catalogue des fourmis d'Europe. *Mittheilungen der Schweizerischen Entomologische Gesellschaft* 5: 441-484.
- Erichson, W.F. 1842. Beitrag zur Insecten-Fauna von Vandiemensland, mitbesonderer Berucksichtigung der geographischen Verbreitung der Insecten. *Archiv fur Naturgeschichte* 8: 83-287.
- Ettershank, G. 1966. A generic revision of the world Myrmicinae related to *Solenopsis* and *Pheidologeton*. *Australian Journal of Zoology* 14: 73-171.
- Fabricius, J.C. 1775. *Systema Entomologiae, sistens insectorum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, and observationibus*: 832 pp. Flensburgi et Lipsiae.
- _____ 1793. *Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, and observationibus* 2: 519 pp. Hafniae.
- _____ 1798. *Supplementum Entomologiae Systematicae*: 572 pp. Hafniae.
- _____ 1804. *Systema Piezatorum*; 439 pp. Brunsvigae.
- Foerster, A. 1850. *Hymenopterologische Studien I. Formicariae*: 74 pp. Aachen.
- Forel, A. 1878. Etudes myrmecologiques en 1878 (premiere partie) avec l' anatomie du gesier des fourmis. *Bulletin de la Societe Vaudoise des Sciences Naturalles* 15: 337-392.
- _____ 1881. Die Ameisen der Antille. *Mittheilungen des Munchener Entomologischen Vereins* 5: 1-16.
- _____ 1886. Indian ants of the Indian Museum, Calcutta. No. 2. *Journal of the Asiatic Society of Bengal. Part 2. Natural science* 55: 239-249.
- _____ 1887. Fourmis recoltees a Madagascar, par le Dr. Conrad Keller. *Mittheilungen der Schweizerischen Entomologische Gesellschaft* 7: 381-389.
- _____ 1890. *Aenictus-Typhlatta* decouverte de M. Wroughton. Nouveaux genres de formicides. *Annales de la Societe Entomologique de Belgique. Comptes-rendus* 34: cii-cxiv.
- _____ 1892a. Quelques fourmis de la fauna mediterraneenne. *Annales de la Societe Entomologique de Belgique* 36: 452-457.
- _____ 1892b. Die Ameisenfauna Bulgariens. (nebst biologischen Beobachtungen.) *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 42: 305-318.
- _____ 1893a. Sur la classification de la famille des formicides, avec remarques synonymiques. *Annales de la Societe Entomologique de Belgique* 37: 161-167.
- _____ 1893b. Les formicides de l'Empire des Indes et de Ceylan. Part 3. *Journal of the Bombay Natural History Society* 8: 17-36.
- _____ 1893c. Note preventive sur un nouveau genre et une nouvelle espece de formicide (campomotide). *Annales de la Societe Entomologique de Belgique* 37: 607-608.
- _____ 1900a. Ponerinae et Dorylinae d'Australie. Recoltes par MM. Turner, Froggatt, Nugent, Chase, Rothney, J.-J. Walker, etc. *Annales*

- de la Societe Entomologique de belgique 44: 54-77.
- _____ 1900b. Les formicides de la l'Empire des Indes et de Ceylan. Part 6. Sous famille Ponerinae. *Journal of the Bombay Natural History Society* 13: 52-65.
- _____ 1900c. Les formicides de la l'Empire des Indes et de Ceylan. Part 7. *Journal of the Bombay Natural History Society* 13: 303-332.
- _____ 1900d. Fourmis de Japon. Nids en toile. *Strongylognathus huberi* et voisins. Fourmiere tribe. *Cyphomyrmex wheeleri*. Fourmis importees. *Mittheilungen der Schweizerischen Entomologische Gesellschaft* 10: 267-287.
- _____ 1901a. Les formicides de la l'Empire des Indes et de Ceylan. Part 8. Sous famille Dorylinae. *Journal of the Bombay Natural History Society* 13: 462-477.
- _____ 1901b. Formiciden des Naturhistorischen Museum zu Hamburg. Neue Calyotomyrmex, Dacryon, Podomyrma, und Echinopla-Arten. *Jahrbuch der Hamburgischen Wssenschaftlichen Anstalten [Mitteilungen aus dem Naturhistorischen Museum]* 18: 45-82.
- _____ 1902. Myrmicinae nouveaux de l'Inde et de Ceylan. *Revue Suisse de Zoologie* 10: 165-249.
- _____ 1905. Ameisen aus Java. Gesammelt von Prof. Karl Kraepelin, 1904. *Jahrbuch der Hamburgischen Wssenschaftlichen Anstalten [Mitteilungen aus dem Naturhistorischen Museum]* 22: 1-26.
- _____ 1911a. Fourmis de Borneo, Singapore, Ceylan, Etc. *Revue Suisse de Zoologie* 19: 23-62.
- _____ 1911b. Ameisen aus Java beobachtet und gesammelt von Herrn Edward Jacobson. 2 Theil. *Notes from the Leyden Museum* 33: 193-218.
- _____ 1911c. Fourmis nouvelles ou interessantes. *Bulletin de la Societe Vaudoise des Sciences Naturelles* 47: 331-400.
- _____ 1911d. Ameisen aus Ceylan, gesammelt von Prof. K. Escherich (einige von Prof. E. Bugnion). In *Escherich, K. Termitenleben auf Ceylon*: 213-228.
- _____ 1912a. H. Sauters Formosa-Ausbeute: Formicidae. *Entomologische Mitteilungen* 1: 45-81.
- _____ 1912b. Descriptions provisoires de genres, sous-genres et especes de formicides des Indes orientales. *Revue Suisse de Zoologie* 20: 761-774.
- _____ 1912c. Einige neue und interessantes Ameisenformen aus Sumatra etc. *Zoologische Jahrbucher Supplement* 15: 51-78.
- _____ 1913. Wissenschaftliche Ergebnisse einer Forschungsreise nach Ostindien, ausgeführt im Auftrage der Kgl. Preuss. Akademie der Wissenschaften zu Berlin von H.V. Buttel-Reepen. 2. Ameison aus Sumatra, Java, Malacca und Ceylon. Gesammelt von Herrn Prof. Dr. v. Buttel-Reepen in den Jahren 1911-1912. *Zoologische Jahrbucher. Abteilung fur Systematik, Geographie und Biologie der Tiere* 36: 1-148.
- Francoeur, A. 1981. Le groupe nearctique *Myrmica lampra*. *Candian Entomologist* 113: 755-759.
- Girard, M. 1879. *Traite elementaire d entomology* 2: 1028 pp. Paris.
- Gotwald, W.H. Jr. 1982. Army Ants (pp. 157-254). In Hermann, H.R. (ed.) *Social Insects* 4: 385 pp. New York.
- Guerin-Meneville, F.E. 1844. Iconographie du regne animal de G. Cuvier, ou representation dapres nature de l'une des especes les plus remarquables et souvent non figurees de chaque genre d'animaux. Insectes. 576 pp.
- Holldobler, B., and E.O.Wilson. 1990. *The Ants*. Cambridge, Mass. 732 pp.
- Jerdon, T.C. 1851. A Catalogue of the species of ants found in southern India. *Mardras Journal of Literature and Science* 17: 103-127.
- Kempf, W.W. 1959. A synopsis of the New World species belonging to the Nesomyrmex-group of the ant genus *Leptothorax* Mayr. *Studia Entomologica* (N.S.) 2: 225-270.
- _____ 1960. *Tranopeltoides* Wheeler, a synonym of *Crematogaster* Lund. *Entomoloical News* 71: 173-175.

- _____ 1967. A synopsis of the Neotropical ants of the genus *Centromyrmex* Mayr. *Studia Entomologica (N.S.)* 9: 401-410.
- _____ 1969. Miscellaneous studies on Neotropical ants. 5. *Studia Entomologica (N.S.)* 12: 273-296.
- _____ 1972. Catalogo abreviado das formigas da Regiao Neotropical. *Studia Entomologica (N.S.)* 15: 3-344.
- Latreille, P.A. 1802. *Histoire Naturelle des Fourmis, et recueil de memoires et dobservations sur les abeilles, les araignees, les faucheurs insectes*. 445 pp. Paris.
- _____ 1804. Tableau methodique des insectes. Classe huitieme. Insectes, Insecta. *Nouveau Dictionnaire d'Histoire Naturelle* 24: 129-200.
- _____ 1810. *Considerations generales sur lordre naturel des animaux composant les classe des Crustaces, des Arachnides, et des Insectes; avec un tableau methodique de leurs genres, disposes en famille*. 444 pp. Paris.
- Lattke, J.E. 1987. Note on the ant genus *Hypoclinea* Mayr, with descriptions of three new species. *Entomologica Scandinavica* 21: 173-178.
- Le Guillou, E.J.F. 1842. Catalogue raisonne des insectes hymenopteres recueillis dans le voyage de circumnavigation des corvettes l'Astrolable et La Zelee. *Annales de la Societe Entomologique de France* 10 (1841): 311-324.
- Linnaeus, C. 1758. *Systema Naturae per renga tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Editio 10. 1: 823 pp. Holmiae.
- _____ 1764. *Museum S:ae R:ae M:tis Ludovicae Ulricaе Reginae Svecorum, Gothorum, Vandalorumque. In que animalia rariora, exotica, imprimis. Insecta and Conchilia describuntur and determinantur. Prodromi instar*: 719 pp. Holmiae.
- Lund, P.W. 1831. Lettre sur les habitudes de quelques fourmis du Bresil, adressee a M. Audouin. *Annales des Sciences Naturelles* 23: 113-138.
- MacKey, W.P. 1993. A review of the New World ant of the genus *Dolichoderus*. *Sociobiology* 22: 1-148.
- Mann, W.M. 1921. The ants of the Fiji Island. *Bulletin of the Museum of Comparative Zoology at Harvard College* 64: 401-499.
- Mayr, G. 1853. Einige neue Ameisen. *Verhandlungen des Zoologisch-Botanischen Vereins in Wien* 2 (1852): 143-150.
- _____ 1855. Formicina austriaca. Beschreibung der bisher im osterreichischen Kaiserataate aufgefundenen Ameisen nebst Hinzufugung jener in Deutschland, in der Schweiz und in Italien vorkommenden Ameisen. *Verhandlungen des Zoologisch-Botanischen Vereins in Wien* 5: 273-478.
- _____ 1861. *Die Europaischen Formiciden*. (Ameisen.): 80 pp. Wien.
- _____ 1862. Myrmecologische Studien. *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 12: 649-776.
- _____ 1863. Formicidarum index synonymicus. *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 13: 385-460.
- _____ 1865. Reise der Osterreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, unter den befehlen des Comodore B. von Wullerstorf-Urbair. *Zoologische Theil. Formicidae*. 119 pp. Wien.
- _____ 1866a. Myrmecologische Beitrage. Sitzungsberichte der k. Akademie der Wissenschaften. *Mathematisch-Naturwissenschaftliche Classe* 53: 484-517.
- _____ 1866b. Diagnosen neuer und wenig gekannter Formiciden. *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 16: 885-908.
- _____ 1867. Adnotationes in monographiam formicidarum Indo-Neerlandicarum. *Tijdschrift voor Entomologie* (2) 2: 33-117.
- _____ 1870. Neue Formiciden. *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 20: 939-996.
- _____ 1872. Formicidae Borneenses collectae a J. Doria et O. Beccari in territorio sarawak

- annis 1865-1867. *Annali del Museo Civico di Storia Naturale di Genova* 2: 133-155.
- _____. 1879. Beitrage zur Ameisen-fauna Asiens. *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 28 (1878): 645-686.
- _____. 1887. Sudamerikanische Formiciden. *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 37: 511-632.
- _____. 1901. Sudamerikanische Formiciden, gesammelt von Dr. Hans Brauns. *Annalen des k.k. Naturhistorischen Hofmuseums Wien* 16: 1-30.
- Menzio, C. 1926. Nuove formiche delle isole Filippine e di Singapore. *Atti della Societa dei Naturalisti e Matematica di Modena* (6) 4: 92-103.
- _____. 1930. Formicidae. (pp.327-332). In Eidmann, H. Entomologische Ergebnisse einer Reise nach Ostasien. *Verhandlungen der k.k. Zoologisch-Botanischen Gesellschaft in Wien* 79: 308-335.
- Moffett, M.W. 1985. Revision of the genus *Myrmoteras*. *Bulletin of the Museum of comparative Zoology* 151: 1-53.
- Nylander, W. 1846. Additamentum adnotationum in monographiam formicarum borealium Europae. *Acta Societatis Scientiarum Fennicae* 2: 1041-1062.
- Olivier, G.A. 1792. Encyclopedie Methodique. Histoire. *Insectes* 6 (part 2):369-704. Paris.
- Perrault, G.H. 1981. *Proceratium deelemani*, nouvelle espece de Kalimantan. *Nouvelle Revue d'Entomologie* 11: 189-193.
- Rigato, F. 1994. Revision of the myrmecine ant genus *Lophomyrmex*, with a review of its taxonomic position. *Systematic Entomology* 19: 47-60.
- Roger, J. 1861. Die Ponera-artigen Ameisen. (schluss.) *Berliner Entomologische Zeitschrift* 5: 1-54.
- _____. 1862. Einige neue exotische Ameisen-Gattungen und Arten. *Berliner Entomologische Zeitschrift* 6: 233-254.
- _____. 1863. Die neue aufgeführten und Arten meines Formiciden-Verzeichnisses, nebst Ergänzung einiger fruher gegeben Beschreibungen. *Berliner Entomologische Zeitschrift* 7: 131-214.
- Santschi, F. 1914. Voyage de Ch.Alluaud et Jeannel en Afrique Orientale (1911-1912). Resultats scientifiques. *Insectes Hymenopteres 2. Formicidae*: 41-148. Paris.
- _____. 1920. Fourmis d'Indo-Chine. *Annales de la Societe Entomologique de Belgique* 60: 158-176.
- _____. 1925. Contribution a la faune myrmecologique de la chine. *Bulletin de la Societe Vaudoise des Sciences Naturelles* 56: 81-96.
- _____. 1927. Revision myrmecologique. *Annales de la Societe Entomologique de Belgique* 67: 240-248.
- Saunders, W.W. 1842. Descriptions of two hymenopterous insects from northern India. *Transactions of the Entomological Society of London* 3 (1841): 57-58.
- Shattuck, S.O. 1992a. Review of the dolichoderine ant genus *Iridomyrmex* Mayr with descriptions of three new genera. *Journal of the Australian Entomological Society* 31: 13-18.
- _____. 1992b. Generic revision of the ant subfamily dolichoderinae. *Sociobiology* 21: 1-181.
- _____. 1993a. Revision of the *Iridomyrmex purpureus* species group. *Invertebrate Taxonomy* 7: 113-149.
- _____. 1993b. Revision of the *Iridomyrmex calvus* species group. *Invertebrate Taxonomy* 7: 1303-1325.
- _____. 1994. Taxonomic Catalog of the ant subfamilies Aneuretinae and Dolichoderinae. *University of California Publications in Entomology* 112: 1-241.
- Shuckard, W.E. 1840. Monograph of the Dorylinae, a family of the Hymenoptera Heterogyna. (Continued from p. 201.) *Annals of Natural History; or Magazine of Zoology, Botany and Geology* 5: 258-271.
- Smith, F. 1851. *List of the specimens of the British animals in the collection of the British Museum* 6. Hymenoptera Aculeata. 134. London.
- _____. 1852. Descriptions of some hymenopterous insects captured in India, with notes on their

- oeconomy, by Ezra T. Downes, Esq., who presented them to the Honourable the East India Company. *Annals and Magazine of Natural History* (2) 9: 44-50.
- _____ 1853. Monograph of the genus *Cryptocerus*, belonging to the group Cryptoceridae-family Myrmiridae-division Hymenoptera Heterogyna. *Transection of the Entomological Society of London* (2) 2: 213-228.
- _____ 1857. Catalogue of the hymenopterous insects collected at Sarawak, Borneo; Mount Ophir; and at Singapore, by A.R. Wallace. *Journal of the Proceedings of the Linnean Society of London, Zoology* 2: 42-88.
- _____ 1858. *Catalogue of the hymenopterous insects in the collection of the British Museum 6 Formicidae*. 216 pp. London.
- _____ 1859. Catalogue of the hymenopterous insects collected, by Mr. A.R. Wallace at the Island of Aru and Key. *Journal of the Proceedings of the Linnean Society of London, Zoology* 3: 132-158.
- _____ 1860a. Description of new species of hymenopterous insects collected, by Wallace at Celebes. *Journal of the Proceedings of the Linnean Society, Zoology* 5 (supplement to volume 4): 57-93.
- _____ 1860b. Catalogue of the hymenopterous insects collected, by Mr. A.R. Wallace in the Island of Bachian, Kaisaa, Amboyna, Gilolo, and at dury in New Guinea. *Journal of the Proceedings of the Linnean Society, Zoology* 5 (supplement to volume 4): 93-143.
- _____ 1861. Catalogue of the hymenopterous insects collected, by Mr. A.R. Wallace in the Island of Ceram, Celebes, Ternate, and Gilolo. *Journal of the Proceedings of the Linnean Society, Zoology* 6: 36-48.
- _____ 1863. Catalogue of the hymenopterous insects collected, by Mr. A.R. Wallace in the Island of Mysol, Ceram, Waigiou, Bouru and Timor. *Journal of the Proceedings of the Linnean Society, Zoology* 7: 6-48.
- _____ 1874. Description of new species of Tenthredinidae, Ichneumonidae, Chrydidae, Formicidae of Japan. *Transection of the Entomological Society of London* (4) 7: 373-409.
- _____ 1877. Description of new species of the genera *Pseudomyrma* and *Tetraponera*, belonging to family Formicidae. *Transection of the Entomological Society of London* (4) 10: 57-72.
- Soulie, J and L.D. Dicko. 1965. La repartition des genres de fourmis de la tribu des Crematogastrini dans la faune ethiopienne et malgache. *Annales de l'Universite d'Abidjan* (Sciences) 1: 85-106.
- Spinola, M. 1851. Compte rendu des Hymenoteres inedits provenants du voyage entomologique de M. Dhiliani dans la Para en 1846. *Extrait des Memoires de l'Academie des Sciences de Turin* (2) 13: 3-78.
- Sykes, W.H. 1835. Description of new species of Indian ants. *Transection of the Entomological Society of London* 1: 99-107.
- Taylor, R.W. 1965. A new species of the ant genus *Dacetonops* from Sarawak. *Breviora* 237: 1-4.
- _____ 1969. The identity of *Dorylozelus mjobergi* Forel. *Journal of Australian Entomological Society* 8: 131-133.
- Taylor, R. W. and D. R. Brown. 1985. *Zoological Catalogue of Australia 2. Hymenoptera, Formicidae, Vespidae and Sphecoidea*: 381 pp. Canberra.
- Terayama, M. and S.Yamane. 1989. Thw army ant genus *Aenictus* from Sumatra with description of three new species. *Japanese Journal of Entomology* 57: 597-603.
- Viehmeyer, H. 1916. Ameisen von Singapore. Beobachtet und gesammelt von H. Overbeck. *Archiv fur Naturgeschichte* 81: 108-168.
- Walker, F. 1859. Characters of some apparently undescribed ceylon insects. *Annals and Magazine of Natural History* 3: 370-376.
- Wang, C. and J. Wu, 1994. Second revisionary studies on genus *Camponotus* Mayr of China. *Journal of Beijing Forestry University (English edition)* 3: 23-34
- Ward, P.S. 2001. Taxonomy, phylogeny and biogeog-

- raphy of the ant genus *Tetraponera* (Hymenoptera: Formicidae) in the Oriental and Australian regions. *Invertebrate Taxonomy* 15: 589-665.
- Watanasit, S., C. Phophuntin and S. Permkam. 2000. Diversity of Ants (Hymenoptera: Formicidae) from Ton Nga Chang Wildlife Sanctuary, Thailand. *Research Article ScieceAsia* 26: 187-194.
- Watanasit, S., Sonthichai and N. Noon-anant. 2003. Preliminary Survey of Ants at Tarutao National Park, Southern Thailand. *Songklanakarin J. Sci. Technol.*, 25 (1): pp. 115-122.
- Weber, N. A. 1944. The Neotropical coccid-tending ants of the genus *Acropygra* Roger. *Annals of Entomological Society of America* 37: 89-122.
- Westwood, J.O. 1835. Hymenopterous Insects, which Mr. Westwood regarded as new to species. *Proceedings of the Zoological Society of London* 3: 86-72.
- _____ 1839. An Introduction to the Modern Classification of Insects; founded on the natural habits and corresponding organisation of the different families 2: (part 11): 193-224.
- _____ 1840. Observations on the genus *Typhlopone*, with descriptions of several exotic species of ants. *Annals and Magazine of Natural History* 6: 81-89.
- Wheeler, W.M. 1909. Ants of Formosa and the Philippines. *Bulletin of the American Museum of Natural History* 26: 333-345.
- _____ 1911. A list of the type species of the genera and subgenera of Formicidae. *Annals of the New York Academy of Sciences* 21: 157-175.
- _____ 1913a. Corrections and additions to list of type species of the genera and subgenera of Formicidae. *Annals of the New York Academy of Sciences* 23: 77-83.
- _____ 1913b. Zoological results of the Abor Expedition, 1911-1912. Hymenoptera 2. Ants (Formicidae). *Records of the Indian Museum* 8: 233-237.
- _____ 1915. The ants of the Baltic Amber. *Schriften der Physikalisch-ökonomischen Gesellschaft zu Königsberg in Pr.* 55: 1-142.
- _____ 1919. The ants of Borneo. *Bulletin of the Museum of Comparative Zoology at Harvard College* 63: 43-147.
- _____ 1922. The ants of the Belgian Congo. *Bulletin of the American Museum of Natural History* 45: 1-1139.
- _____ 1927. Burmese ants collected by Professor G.E. Gates. *Psyche* 34: 42-46.
- _____ 1928a. Ants collected by Professor F. Silvestri in China. *Bolletino del Laboratorio di Zoologia generale e agraria del R. Istituto Superiore agrario di Portici* 22: 3-38.
- _____ 1928b. Ants collected by Professor F. Silvestri Japan and Korea. *Bolletino del Laboratorio di Zoologia generale e agraria del R. Istituto Superiore agrario di Portici* 22: 96-125.
- _____ 1928c. A new species of *Probolomyrmex* from Java. *Psyche* 35: 7-9.
- _____ 1929. A *Camponotus mermithergate* from Argentina. *Psyche* 36: 102-106.
- Wheeler, W.M. and J.W. Chapman. 1925. The ants of the Philippine Islands. Part 1. Dorylinae and Ponerinae. *Philippine Journal of Science* 28: 49-73.
- Wilson, E.O. 1955. The status of the ant genus *Microbolbos* Donisthorpe. *Psyche* 62: 136.
- _____ 1958. Studies on the ant fauna of Melanesia. Part 1 The tribe Leptogenyini. 2. The tribe Amblyoponini. *Bulletin of the Museum of Comparative Zoology at Harvard College* 118: 101-153.
- _____ 1964. The true army ants of the Indo-Australian area. *Pacific Insects* 6: 427-483.
- Wiwatwitaya, D. and W. Jaitrong. 2001. *Identification Guide to the Ant Genera of Khao Yai National Park*. Kasetsart University. 110 pp.
- Xu, Z. 2000. Two new genera of the ant subfamilies Dorylinae and Ponerinae (Hymenoptera: Formicidae) From China. *Zoological Research* (4): 297-302.

Received: 26 August 2004
Accepted: 1 September 2004