

Marine Algae of the Eastern Caraga Region, Philippines, I. Historical Survey and an Annotated List of the Brown Algae (Phaeophyceae)

Dan Anthony U. Bataan^{1,5}, Cynthia P. Sajot², Paul John L. Geraldino³ and Lawrence M. Liao^{4,*}

¹ *Surigao State College of Technology, Main Campus, Surigao City, Philippines*

² *Surigao del Sur State University, Lianga Campus, Lianga, Surigao del Sur, Philippines*

³ *Department of Biology, University of San Carlos, Cebu City, Philippines*

⁴ *Graduate School of Integrated Sciences for Life, Hiroshima University, Higashi-Hiroshima, Japan*

⁵ *Present address: Kuroshio Science Program, Graduate School of Integrated Arts and Sciences, Kochi University, Nankoku, Japan*

* Corresponding author: lliao@hiroshima-u.ac.jp

ABSTRACT

The marine algal flora of the Philippines is highly diverse owing to its strategic location in the middle of the vast Indo-Pacific biogeographic region. However, the diversity of marine algae in many parts of the country remains unknown due to geographic isolation that limits botanical exploration. The Caraga region in northeastern Mindanao Island is one such poorly known locality. This study was aimed to produce a consolidated list of published records of the brown algae from the region, including a historical survey of the few studies conducted there previously. A total of 32 species, 11 genera, three families and three orders of marine brown algae were recorded. Information revealed some economically-important species which could be tapped for livelihood creation in coastal villages there. Baseline information presented in this study will be useful for resource exploitation and policy formulation.

Keywords: botanical inventory, brown algae, Mindanao, phycology.

INTRODUCTION

The political region of Caraga in the Philippines is composed of five provinces located on the northeastern part of the island of Mindanao including several large offshore islands. Its eastern coast faces the vast Pacific Ocean from where the warm equatorial currents bring propagules that contribute to the rich marine biodiversity found there. In addition, several typhoons hit the area often bringing storm surges that cause greater water circulation. Some upwelling from the nearby Philippine Trench could also contribute to the high marine productivity seen there.

Compared to other regions in the Philippines, the Caraga region is poorly known in terms of its biodiversity despite the variety of favorable geographical features found there, i.e., the long Agusan River and the vast Agusan Marsh, the majestic Diwata mountain range, the Lake Mainit basin and the several offshore islands with their numerous lagoons and rugged coastlines. During the American colonial period in the early 20th century which saw the start of biological exploration and collecting mainly in the Christianized parts of the Philippines, the Caraga region was among those largely excluded. Instead, the attention of the colonizers was

focused on the development of gold and nickel mining activities in the eastern part of the region which continue to this day at immeasurable environmental and social costs.

Limited biological exploration of the Caraga region started during the early years of the American colonial period. Among the pioneers was Edward H. Taylor who was an American herpetologist at that time who explored the region and published on many species known to science for the first time (Borrell, 2013; Taylor, 1920). Among the earliest accounts of marine fauna there were based on the collections of Francisco Nemenzo, Sr. (formerly of the University of the Philippines Diliman) and Exuperancio A. Montecillo (formerly of the University of San Carlos) who led a team of biologists which found new species of scleractinian corals from the waters of Aras-asan Island in Surigao del Sur in 1977 and published subsequently by Nemenzo and Montecillo (1981).

Ernani G. Meñez, who at that time was a research assistant and graduate student at the University of Hawai'i studying under Maxwell S. Doty appears to be the first to collect seaweed samples from the eastern Caraga region. He first collected from Barangay Mangagoy,

Bislig, Surigao del Sur on 29 June 1958, and stayed in the general vicinity for about one week and collected around Mawes Island in Hinatuan and Sanco Point in Maribojoc in Surigao del Sur. His collections currently deposited at the herbarium of the Bernice P. Bishop Museum in Honolulu were studied and published by Taylor (1964, 1977); Hollenberg (1968, 1968a); Gilbert and Doty (1969); Saito (1969); Xia and Abbott (1987); Liao (1990).

Gregorio T. Velasquez, professor of botany at the University of the Philippines Diliman at that time, collected some species of the marine red algal genus *Galaxaura* from Loreto in Dinagat Island on May 1965 and which were published by Cordero (1977). Later, visits to various sites in the Surigao provinces by the team from the National Museum in Manila led by Paciente A. Cordero, Jr. in the 1970s and 1980s produced many samples which were published by Cordero (1980, 1980a); Cadano and Trono (1987); Modelo and Umezaki (1995); Liao *et al.* (1997), among others. Based on samples collected by his students from various parts of the Surigao provinces, Trono (1997, 2004, 2017, 2017a, 2018) added to the growing species list of marine algae there. He also published numerous species from various sites in the Caraga region collected during the Pacific Seaboard Expedition from 2000 to 2003 (Trono 2017a, 2018).

More recently, Stefano G.A. Draisma, a research fellow at the University of Malaya in Kuala Lumpur at that time made extensive collections of marine algae around the Philippines. From 10-12 April 2014, he collected samples from Siargao Island. Among those he collected were samples of the new species, *Halymenia malaysiana* P.-L. Tan, P.-E. Lim, S.-M. Lin and S.-M. Phang which was formally described by Tan *et al.* (2015).

Home-grown biologists also contributed to the study of marine algae in subsequent years. Cynthia P. Sajot of the Surigao del Sur State College at that time conducted a comprehensive survey of the marine algal flora of several sites around Lianga Bay in Surigao del Sur (Sajot, 2006). Fajardo *et al.* (2016) at the Caraga State University surveyed the marine algae and seagrasses near the mining site in Barangay Cagdianao, Placer, Surigao del Norte.

This paper attempted to produce a comprehensive unified list of all species of marine brown benthic algae (Phaeophyceae) reported thus far from the eastern Caraga region, namely the provinces of Surigao del Norte, Surigao del Sur and the Dinagat Islands to serve as baseline information for purposes of resource

management and exploitation, environmental protection, biodiversity conservation and the formulation and implementation of appropriate policies and legislation thereof. This study also aimed to encourage further studies of the marine algae in the Caraga region, to sample in areas where no data are available, to identify species of economic and ecological importance, among other goals.

MATERIALS AND METHODS

This review is based entirely on a comprehensive survey of the published taxonomic literature of marine algae from the three coastal provinces of eastern Caraga region, comprising of Surigao del Norte, Surigao del Sur and the Dinagat Islands. Sources include online materials, library collections and materials compiled personally by the authors. Records were retrieved from each reference and compiled into the checklist presented in this paper. Actual samples were not examined in this survey, instead critical annotations are provided for species whose identity are either doubtful or questionable on biogeographical and other grounds. Data-mined names and records were listed and arranged following the classification system of Dawes and Mathieson (2008) which was selected arbitrarily for adoption in this study. Validity of each names were verified by consulting the latest taxonomic literature and by referring to AlgaeBase (Guiry and Guiry 2020).

RESULTS AND DISCUSSION

The list of all known published species of marine benthic brown algae from the eastern Caraga region, namely the provinces of Surigao del Norte, Surigao del Sur and the Dinagat Islands revealed a total of 32 species, 11 genera, three families and three orders. The list is presented below following the classification scheme of Dawes and Mathieson (2008). Sources of original reports are indicated inside brackets.

HETEROKONTOPHYTA
Class PHAEOPHYCEAE
Order DICTYOTALES
Family DICTYOTACEAE

***Canistrocarpus cervicornis* (Kützing) De Paula and De Clerck** [Trono, 2018]

***Dictyopteris jamaicensis* Taylor** [Fajardo *et al.*, 2016]

Remarks: *D. jamaicensis* is a tall species (15 cm or more) first described from the deep waters off Jamaica by Taylor (1960) and is likely confined to the waters of the Caribbean. The name has become of convenient use

as applied to larger (>10 cm) plants collected from the Philippines (Silva *et al.*, 1987). Obviously Philippine representatives of this genus are in need of critical study.

***Dictyopteris repens* (Okamura) Børgesen** [Trono, 2018]

***Dictyota dichotoma* (Hudson) J.V. Lamouroux** [Trono, 2018]

Remarks: This record should be treated with caution as this species seems restricted to European waters. Kraft (2009: 164) opined that species lacking structural or habit peculiarities, except perhaps for having evenly dichotomous branching, have been conveniently placed under this species.

***Dictyota mertensii* (Martius) Kützinger** [Trono, 2018]

***Lobophora variegata* (J.V.Lamouroux) Womersley ex E.C. Oliveira** [Trono, 2018]

Remarks: This identification should be considered tentative as molecular studies have revealed a bewildering number of cryptic species therein (see Vieira *et al.*, 2016).

***Padina australis* Hauck** [Geraldino *et al.*, 2005; Sajot, 2006; Trono, 2018]

***Padina boryana* Thivy** [Geraldino *et al.*, 2005]

***Padina fernandeziana* Skottsberg and Levring** [Geraldino *et al.*, 2005]

Remarks: The singular report of this apparently endemic Eastern Pacific species needs to be reevaluated in the light of many new species recently described from the Indo-West Pacific (Ni-Ni Win *et al.*, 2010, 2011, 2012).

***Padina japonica* Yamada** [Trono, 2018]

***Padina minor* Yamada** [Geraldino *et al.*, 2005; Sajot, 2006; Fajardo *et al.*, 2016; Trono, 2018]

Remarks: Philippine records of this species need to be treated with caution as there is a tendency among researchers to apply this name indiscriminately to *Padina* species which are “smaller than usual” disregarding other plant features (Modelo and Umezaki, 1884; Geraldino *et al.*, 2005; Tsutsui *et al.*, 2005)

***Padina moffittiana* Abbott and Huisman** [Geraldino *et al.*, 2005]

Remarks: Originally described from Hawaii, this species has now been reported from southern Japan (Ni-Ni-Win *et al.*, 2010) and several sites in the Philippines

(Geraldino *et al.*, 2005).

***Padina sanctae-crucis* Børgesen** [Geraldino *et al.*, 2005]

***Padina* sp.** [Fajardo *et al.*, 2016]

Order ECTOCARPALES
Family SCYTOSIPHONACEAE

***Colpomenia sinuosa* (Mertens ex Roth) Derbés and Solier ex Castagne** [Trono, 2018; Sajot, 2006, as *Colpomenia* sp.]

Remarks: The record of Sajot (2006) most likely refers to the amorphous and convoluted sac-like species listed above.

***Hydroclathrus clathratus* (C.Agardh) Howe ex Britton and Millsbaugh** [Trono, 2018; Sajot, 2006, as *Hydroclathrus* sp.]

Remarks: The identity of the sample of Sajot (2006) cannot be ascertained without a critical examination of voucher specimens. In the Philippines, two reticulate species have been customarily reported: *H. clathratus* which is a cosmopolitan species characterized by branches of irregular diameters (>3 mm), and *H. tenuis* Tseng and Lu, a more fragile species with consistently finer branches (<2 mm) and a more restricted distribution than the former. Kraft and Abbott (2003), Santiañez and Wynne (2019), and Santiañez *et al.* (2017, 2018) have identified superficially similar reticulate species as new species and new genera with the aid of molecular and morphological evidence.

Order FUCALES
Family SARGASSACEAE

***Hormophysa cuneiformis* (Gmelin) Silva** [Liao *et al.*, 1997; Trono, 2018; Sajot, 2006 as *Hormophysa* sp.]

Remarks: The triquetrous nature of the thallus is distinctive for this species. In mature specimens, air vesicles are embedded into the thallus. Formerly placed within the Cystoseiraceae, this genus is now classified within the Sargassaceae after Draisma *et al.* (2010) synonymized the two families. The unidentified species reported by Sajot (2006) is almost certainly *H. cuneiformis*.

***Sargassum aquifolium* (Turner) J.Agardh**
=*Sargassum heterocystum* Montagne [Cordero, 1980; Modelo and Umezaki, 1995]

***Sargassum ilicifolium* (Turner) C.Agardh**
=*Sargassum cristaefolium* C.Agardh [Trono,

1997; Fajardo *et al.*, 2016; Trono, 2018]
 =*Sargassum duplicatum* J. Agardh [Cordero, 1980, as *Sargassum* sp. 1, 1980a]

Remarks: The duplicated margins consisting of serrated or dentate rows may be seen either disposed entirely on leaf margins or only on the distal half of the leaf.

***Sargassum grandulifolium* Grunow ex Piccone** [Modelo and Umezaki, 1995]

***Sargassum oligocystum* Montagne** [Trono, 1997]

***Sargassum paniculatum* J. Agardh** [Fajardo *et al.*, 2016]

***Sargassum polycystum* C. Agardh** [Trono 1997; Fajardo *et al.*, 2016]

=*Sargassum myriocystum* J. Agardh [Modelo and Umezaki, 1995]

Remarks: This species is among the easiest to identify with its distinctively ‘spiny’ main axis which are, in fact, beset with cryptostomata that have expanded ostioles (Chiang *et al.*, 1992).

***Sargassum samarense* Trono** [Trono, 2018]

***Sargassum siliquosum* J. Agardh** [Trono, 2018]

***Sargassum sullivanii* Trono** [Trono, 2018]

***Sargassum yoshidae* Trono** [Trono, 2018]

***Sargassum* sp.** [Sajot, 2006; Fajardo *et al.*, 2016]

***Sirophysalis trinodis* (Forsskål) Kützing**

=*Cystoseira trinodis* (Forsskål) C. Agardh [Liao *et al.*, 1997]

Remarks: For so many years, this peculiar species was recognized under *Cystoseira* with a distinctively Indo-Pacific distribution quite different from its congeners that are restricted within the Mediterranean basin, its presumed center of distribution. Draisma *et al.* (2010) removed it from *Cystoseira* and transferred it into its own genus and at the same time subsuming the Cystoseiraceae (containing *Cystoseira*, *Hormophysa* and *Sirophysalis*) under the Sargassaceae based on molecular evidence.

***Turbinaria condensata* Sonder** [Taylor, 1964]

Turbinaria conoides* (J. Agardh) Kützing f. *conoides [Trono, 2018]

***Turbinaria conoides* f. *laticuspidata* Taylor** [Taylor, 1964]

***Turbinaria decurrens* Bory** [Taylor, 1964; Trono, 1997, 2018]

Remarks: The three-sided obpyramidal shape of the main photosynthetic blade is distinctive. Another morpho-logically similar species is *T. murrayana* Barton which is delineated from *T. decurrens* by “having short, thick, unbranched stem,...” among other features.

***Turbinaria ornata* (Turner) J. Agardh** [Trono, 2018]

Records of marine benthic algae in the Philippines are largely scattered due to various factors. Geographic distance and isolation of some habitats can be cited for the uneven data thus far available, especially in an archipelagic country like the Philippines where safe access is of vital concern. Weather concerns are also taken into account. In the study area of the eastern Caraga region and throughout the Pacific coasts of the Philippines (Evangelista *et al.*, 2015), typhoons pose serious threats to collecting activities. Another important factor to explain the patchy distributional records of marine algae in the Philippines is the lack of coordinated collecting efforts especially in far flung regions like Caraga. For practical reasons, collections have focused on easily accessible areas and those located near research centers in the national capital.

This survey revealed the obvious absence of common tropical marine floral elements. For instance, some unaccounted for in the literature survey were common epiphytic brown algae such as those belonging to uniseriate *Hincksia* and *Feldmannia* (Acinetosporaceae) and multiseriate *Sphacelaria* (Sphacelariaceae). Silva *et al.* (1987) reported 3-4 species of each of these genera from the Philippines where they form dense epiphytic tufts on larger algae, but it is believed the diversity of these microscopic brown species are understated, at least in the Philippines. Another missing element are the crustose brown algae belonging to the Ralfsiales.

Among the most common species listed are members of *Sargassum* which are an emerging source of alginates (Fawzy *et al.*, 2017) in many parts of the world. Alginate extraction residues have also been used to absorb toxic heavy metals (Cardoso *et al.*, 2017). Many species are also of biomedical importance for their cytotoxic (Shofia *et al.*, 2018) and anti-inflammatory effects (Luo *et al.*, 2019). In addition to *Sargassum*, biomass from species of *Padina* have been used as potential agents for aquatic environmental remediation (Sheng *et al.*, 2004).

The marine flora of the eastern Caraga region may be

considered moderately rich as evidenced by the number and diversity of species recorded there. The absence of some common tropical species in the literature survey suggested that more surveys need to be conducted in eastern Caraga to account for species that are commonly expected in tropical marine ecosystems. To account for species that may exhibit seasonality, sampling should be done during the dry and wet seasons of the year. Notwithstanding the gaps exposed in this survey, this compilation of past records is an important step towards completing the total phytogeographical picture of the Philippines. Similar compilations of records have helped immensely in consolidating scattered records (Silva *et al.*, 1987) and the updating and revision of records have kept up with changes in taxonomy (Liao and Young, 2002; Evangelista *et al.*, 2015). Without doubt, the records of the marine flora of the Caraga region will increase with more collections through time. The information in this report will hopefully point out a number of taxonomic gaps. As succinctly stated by Silva *et al.* (1987), works such as species surveys and listings “are not intended to endure like monoliths,” but their “rate of obsolescence [becomes] an index of progress” for the study of marine algae including those to be found in the Caraga region.

CONCLUSION

This inventory of the species of marine algae, specifically of the brown algal species (Phaeophyceae), of the eastern Caraga region represents an attempt to compile information which would otherwise be obscured or lost. The current effort is important considering that whatever widely scattered available information needs verification. This effort is important as there is a general lack of research cooperation and coordination, sporadic publication in the gray literature, scarcity of taxonomic expertise, and low emphasis on the importance of seaweed resources, among other reasons. The current survey is a contribution towards increasing the profile of seaweed resources for resource managers and policy makers especially in an area like the Caraga region whose marine resources are obviously plentiful but only minimally exploited. Annotated checklists like the current work can serve to highlight the biodiversity of marine algae, identify those with potential economic importance which can be tapped for economic improvement for coastal communities. Information herein can also be used to help technology transfer by extension workers for income generation, among many other applications.

ACKNOWLEDGMENTS

A travel grant was provided to LML by the Graduate School of Integrated Sciences for Life of Hiroshima University. The staff at the Hiroshima University Library provided help in locating some references used in this study. Some hard-to-find literature were also provided by some colleagues for which the authors are grateful. The authors thank Dr. Suttikarn Sutti of the Thailand Natural History Museum and an anonymous reviewer for providing valuable insights useful in the revision of this manuscript.

REFERENCES

- Borrell, B. 2013. The spy who loved frogs. *Nature* 501: 150–153.
- Cadano, M.A. and G.C. Trono. 1987. The genus *Jania* (Cryptonemiales, Rhodophyta) in the Philippines. *Philippine Journal of Science* 116: 99–117.
- Cardoso, S.L., C.S.D. Costa, E. Nishikawa, M.G.C. da Silva and M.G.A. Vieira. 2017. Biosorption of toxic metals using the alginate extraction residue from the brown algae *Sargassum filipendula* as a natural ion-exchanger. *Journal of Cleaner Production* 165: 491–499.
- Chiang, Y.M., T. Yoshida, T. Ajisaka, G.C. Trono, C.K. Tseng and B. Lu. 1992. Distribution and variation in *Sargassum polycystum* C. Agardh (Fucales, Phaeophyta). In: I.A. Abbott (ed.), *Taxonomy of Economic Seaweeds with Reference to Some Pacific and Western Atlantic Species*. California Sea Grant College Program, La Jolla, USA, pp. 93–106.
- Cordero, P.A. 1977. Studies on Philippine marine red algae. *Special Publications from the Seto Marine Biological Laboratory* 4: 1–258.
- Cordero, P.A. 1980. Taxonomy and distribution of Philippine useful seaweeds. *National Research Council of the Philippines Bulletin* 81: 1–78.
- Cordero, P.A. 1980a. Phycological observations VII. On the occurrence and phycogeographical distribution of *Sargassum duplicatum* and *Halicoryne wrightii* in the Philippines. *Publications of the Seto Marine Biological Laboratory* 25: 27–38.
- Dawes, C.J. and A.C. Mathieson. 2008. *The Seaweeds of Florida*. University Press of Florida, Gainesville, USA. 656 pp.
- Draisma, S.G.A., E. Ballesteros, F. Rosseau and T. Thibaut. 2010. DNA sequence data demonstrate the polyphyly of the genus *Cystoseira* and other Sargassaceae genera (Phaeophyceae). *Journal of Phycology* 46: 1329–1345.
- Evangelista, E.V., P.A. Cordero, L.T. Evangelista and L.M. Liao. 2015. Macrobenthic algae of the southern coasts of Catanduanes Island, Philippines: historical account and critical review of records. *The Thailand*

- Natural History Museum Journal* 9: 95–108.
- Fajardo, D.R.M., R.P. Magdugo and R.A. Seronay. 2016. Diversity and taxonomic assessment of marine benthic macrophytes near mining situated area in Cagdianao. Claver, Surigao del Norte, Philippines. *Journal of Biology and Environmental Science* 8: 49–56.
- Fawzy, M.A., M. Gomaa, A.F. Hifney and K.M. Abdel-Gawad. 2017. Optimization of alginate alkaline extraction technology from *Sargassum latifolium* and its potential antioxidant and emulsifying properties. *Carbohydrate Polymers* 157: 1903–1912.
- Geraldino, P.J.L., L.M. Liao and S.M. Boo. 2005. Morphological study of the marine algal genus *Padina* (Dictyotales, Phaeophyceae) from southern Philippines: 3 species new to Philippines. *Algae* 20: 99–112.
- Gilbert, W.J. and M.S. Doty. 1969. Some additional records of Philippine marine Chlorophyta. *Micronesica* 5: 121–130.
- Guiry, M.D. and G.M. Guiry. 2020. AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. <http://www.algaebase.org>; searched on 11 February 2020.
- Hollenberg, G.J. 1968. An account of the species of *Polysiphonia* of the central and western tropical Pacific Ocean, I. Oligosiphonia. *Pacific Science* 22: 56–98.
- Hollenberg, G.J. 1968a. An account of the species of the red alga *Herposiphonia* occurring in the central and western tropical Pacific Ocean. *Pacific Science* 22: 536–559.
- Kraft, G.T. 2009. *Algae of Australia: Marine Benthic Algae of Lord Howe Island and the Southern Great Barrier Reef, 2. Brown Algae*. Australian Biological Resources Study, Canberra and CSIRO Publishing, Melbourne. 364 pp.
- Kraft, G.T. and I.A. Abbott. 2003. *Hydroclathrus* (Scytosiphonaceae, Phaeophyceae): conspectus of the genus and the proposal of new species from Australia and Hawaii. *Phycological Research* 51: 244–258.
- Liao, L.M. 1990. Discovery of the “*Falkenbergia*” phase of *Asparagopsis taxiformis* (Rhodophyta: Bonnemaisoniales) in Philippine waters. *The Philippine Scientist* 27: 5–10.
- Liao, L.M., R.B. Modelo and I. Umezaki. 1997. Taxonomy and phytogeography of Philippine Cystoseiraceae (Phaeophyta: Fucales). *The Philippine Scientist* 34: 21–32.
- Liao, L.M. and J.G. Young. 2002. Marine algae of the Sulu Sea islands, Philippines, I: Introduction, historical account and additional records from the Tubbataha reefs. *The Philippine Scientist* 39: 15–35.
- Luo, D., Z. Wang and K. Nie. 2019. Structural characterization of a novel polysaccharide from *Sargassum thunbergii* and its antioxidant and anti-inflammation effects. *PLoS One* 14(10): e0223198.
- Modelo, R.B. and I. Umezaki. 1984. *Padina* and *Pocockiella* species of Luzon Is., Philippines. *Memoirs of the College of Agriculture, Kyoto University* 125: 11–34.
- Modelo, R.B. and I. Umezaki. 1995. Contribution to the study of the genus *Sargassum* (Fucales, Phaeophyceae) of the Philippines. *Philippine Journal of Science, Special Issue*, 1–50.
- Modelo, R.B., I. Umezaki and L.M. Liao. 1998. Morphology of *Sargassum corderoi* (Phaeophyta, Fucales), a new seaweed species from central Philippines. *Algae, Korean Journal of Phycology* 13: 79–83.
- Nemenzo, F. and E. Montecillo. 1981. Four new scleractinian species from Arangasa Islet (Surigao del Sur province, Philippines). *The Philippine Scientist* 18: 120–128.
- Ni-Ni-Win, T. Hanyuda, S. Arai, M. Uchimura, A. Prathep, S.G.A. Draisma, S.M. Phang, I.A. Abbott, A.J.K. Millar and H. Kawai. 2011. A taxonomic study of the genus *Padina* (Dictyotales, Phaeophyceae) including the descriptions of four new species from Japan, Hawaii and the Andaman Sea. *Journal of Phycology* 47: 1193–1209.
- Ni-Ni-Win, T. Hanyuda, S. Arai, M. Uchimura, A. Prathep, S.G.A. Draisma, Soe-Htun and H. Kawai. 2010. Four new species of *Padina* (Dictyotales, Phaeophyceae) from the western Pacific Ocean and reinstatement of *Padina japonica*. *Phycologia* 49: 136–153.
- Ni-Ni-Win, T. Hanyuda, S.G.A. Draisma, E. Verheij, W.F. Prud'homme van Reine, P.-E. Lim, S.-M. Phang and H. Kawai. 2012. Morphological and molecular evidence for two new species of *Padina* (Dictyotales, Phaeophyceae), *P. sulcata* and *P. calcarea*, from the central Indo-Pacific region. *Phycologia* 51: 576–585.
- Saito, Y. 1969. The algal genus *Laurencia* from the Hawaiian Islands, the Philippine Islands and adjacent areas. *Pacific Science* 23: 148–160.
- Sajot, C.P. 2006. Assessment of macrobenthic algae in Lianga Bay, eastern Mindanao, Philippines. *Surigao del Sur Polytechnic State College Research, Development and Extension Journal* 2: 91–98.
- Santiañez, W.J.E., K.M. Lee, S. Uwai, A. Kurihara, P.J.L. Geraldino, E.T. Ganzon-Fortes, S.M. Boo and K. Kogame. 2017. Untangling nets: elucidating the diversity and phylogeny of the clathrate algal genus *Hydroclathrus*, with the description of a new genus *Tronoella* (Scytosiphonaceae, Phaeophyceae). *Phycologia* 57: 61–78.

- Santiañez, W.J.E., E.C. Macaya, K.M. Lee, G.Y. Cho, S.M. Boo and K. Kogame. 2018. Taxonomic assessment of the Indo-Pacific Scytosiphonaceae (Phaeophyceae): *Hydroclathrus rapanuii* sp. nov. and *Chnoospora minima* from Easter Island, with proposal of *Dactylosiphon* gen. nov. and *Pseudochnoospora* gen. nov. *Botanica Marina* 61: 47–64.
- Santiañez, W.J.E. and M.J. Wynne. 2019. Evidence for the treatment of *Talarodictyon tilesii* as an older taxonomic synonym of *Hydroclathrus stephanosorus* (Scytosiphonales, Phaeophyceae). *Phycological Research* 67: 82–85.
- Sheng, P.X., L.H. Tan, J.P. Chen and Y.-P. Ting. 2004. Biosorption performance of two brown marine algae for removal of chromium and cadmium. *Journal of Dispersion Science and Technology* 25: 681–688.
- Shofia, S.I., K. Jayakumar, A. Mukherjee and N. Chandrasekaran. 2018. Efficiency of brown seaweed (*Sargassum longifolium*) polysaccharides encapsulated in nanoemulsion and nanostructured lipid carrier against colon cancer cell lines HCT 116. *RSC Advances* 8: 15973–15984.
- Silva, P.C., E.G. Meñez and R.L. Moe. 1987. Catalog of the marine benthic algae of the Philippines. *Smithsonian Contributions to the Marine Sciences* 27: 1–179.
- Tan, P.-L., P.-E. Lim, S.-M. Lin, S.-M. Phang, S.G.A. Draisma and L.M. Liao. 2015. Foliose *Halymenia* species (Halymeniaceae, Rhodophyta) from Southeast Asia, including a new species, *Halymenia malaysiana* sp. nov. *Botanica Marina* 58: 203–217.
- Taylor, E.H. 1920. Philippine amphibia. *Philippine Journal of Science* 16: 213–359.
- Taylor, W.R. 1960. *Marine Algae of the Eastern Tropical and Subtropical Coasts of the Americas*. The University of Michigan Press, Ann Arbor, USA, 870 pp.
- Taylor, W.R. 1964. The genus *Turbinaria* in eastern seas. *Journal of the Linnean Society of London, Botany* 58: 475–490.
- Taylor, W.R. 1977. Notes on plants of the genus *Caulerpa* in the herbarium of Maxwell S. Doty at the University of Hawaii. *Atoll Research Bulletin* 208: 1–17.
- Trono, G.C. 1997. *Field Guide & Atlas of the Seaweed Resources of the Philippines*. Bookmark Inc., Makati City, Philippines, 306 pp.
- Trono, G.C. 2004. *Field Guide & Atlas of the Seaweed Resources of the Philippines, Volume 2*. Bureau of Agricultural Research, Quezon City, Philippines, 260 pp.
- Trono, G.C. 2017. *Field Guide & Atlas of the Seaweed Resources of the Philippines: Epiphytes & Microbenthic Algae, Volume III*. Bureau of Agricultural Research, Quezon City, Philippines, 230 pp.
- Trono, G.C. 2017a. *An Illustrated Seaweed Flora Collected Along the Pacific Seaboard of the Philippines, Part I. Cyanophyceae and Rhodophyceae*. National Academy of Science and Technology, Taguig City, Philippines, 168 pp.
- Trono, G.C. 2018. *An Illustrated Seaweed Flora Collected Along the Pacific Seaboard of the Philippines, Part II. Chlorophyceae and Phaeophyceae*. National Academy of Science and Technology, Taguig City, Philippines, 166 pp.
- Tsutsui, I., Q.N. Huynh, H.D. Nguyen, S. Arai and T. Yoshida. 2005. *The Common Marine Plants of Southern Vietnam*. 2nd ed. Japan Seaweed Association, Tosa. 250 pp.
- Vieira, C., O. Camacho, M.J. Wynne, L. Mattio, R.J. Anderson, J.J. Bolton, M. Sansón, S. D'hondt, F. Leliaert, S. Fredericq, C. Payri and O. De Clerck. 2016. Shedding new light on old algae: Matching names and sequences in the brown algal genus *Lobophora* (Dictyotales, Phaeophyceae). *Taxon* 65: 689–707.
- Xia, B. and I.A. Abbott. 1987. New species of *Polycavernosa* Chang & Xia (Gracilariaceae, Rhodophyta) from the western Pacific. *Phycologia* 26: 405–418.

Received: 27 July 2020

Accepted: 9 December 2020

Published Online: 29 April 2021

Published in Print: 30 June 2021